

Original Article: Determining the Contribution of Hyperlipidemia to Mortality Post anesthesia in Patients who are Candidates for Coronary Artery Graft Surgery

Khosrow Hashemzadeh¹, Marjan Dehdilan^{2*}

1. Department of Heart Surgery, Tabriz University of Medical Sciences, Tabriz, Iran. (ORCID: 0000-0001-5551-906X)
2. Department of Anesthesiology, Tuberculosis and Lung Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran (ORCID: 0000-0001-8119-3198)



Citation K. Hashemzadeh, M. Dehdilan, **Investigating the Effectiveness of Ketorolac in Controlling Pain after Abdominal Surgeries**, *EJCMPR*. 2023; 2(2):159-168.

 <https://doi.org/10.5281/zenodo.8004716>

Article info:

Received: 01 May 2023

Accepted: 06 Jun 2023

Available Online:

ID: EJCMPR-2306-1043

Checked for Plagiarism: Yes

Peer Reviewers Approved by:

Dr. Amir Samimi

Editor who Approved Publication:

Dr. Frank Rebout

Keywords:

Hyperlipidemia, Mortality, Post-Anesthesia, CABG

ABSTRACT

Introduction: Statins are powerful lipid-lowering drugs that have been shown to be effective in preventing heart disease and reducing the risk of death and heart attack. It is unclear whether hyperlipidemic patients undergoing coronary artery bypass grafting benefit from the lipid-lowering effects of statins. We sought to determine whether prior statin therapy could affect the outcome of hyperlipidemic patients undergoing coronary artery bypass grafting. **Material and Methods:** In this cross-sectional descriptive study conducted between 2012 and 2015, hyperlipidemia to mortality post anesthesia in patients who are candidates for coronary artery graft surgery was investigated. **Results:** Risk-adjusted multivariate logistic regression analysis showed that statin-treated hyperlipidemic (odds ratio, 0.42; 95% confidence interval, 0.26-0.69; P = .0007) ratio was 0.42; confidence interval, 0.26-0.69; P = .0007) was independently associated with a reduction in major in-hospital cardiac events, but not in-hospital mortality. A similarity score based on the previous 14 risk factors was performed to further control for bias. After similar correlations, randomized controlled trials confirmed that statin-treated hyperlipidemia and non-statin-treated eulipidemia were associated with reductions in major cardiovascular hospitalizations (difference odds ratio, 0.41; 95% confidence, 0.24--0.71 [P. .0013] and odds) rate is 0. **Conclusion:** Although there was no increase in MACE in these normolipidemic patients who did not receive prior statin therapy, results from other CABG studies 6,7,8,9,16 and 25 in patients with heart disease were background in all patients who received CABG. LDL-C levels may benefit from long-term statin therapy. Although some of these patients did not see immediate short-term benefit, this study did not examine the long-term outcomes and future cardiovascular events of MACE. This group of patients may also benefit from a reduction in long-term MACE with statin therapy despite low preoperative LDL-C levels.

*Corresponding Author: Fariborz Roustaa (mrjandehdilani@gmail.com)

Introduction

Hyperlipidemia is considered to be one of the main risk factors associated with vascular endothelial damage leading to atherosclerotic plaque formation and coronary artery disease (CAD) leading to recurrent ischemic cardiovascular events. 3-Hydroxy-3-methylglutaryl-CoA reductase inhibitors, statins, are considered potent lipid-lowering agents and have been shown to be effective in primary and secondary prevention of CAD by lowering lipoprotein cholesterol (LDL-C) levels. To reduce the risk of death and adverse cardiovascular events in patients with hyperlipidemia and normcholesterolemia Statin therapy as a surgical intervention has been shown to reduce heart disease and the risk of premature death after heart surgery, suggesting a beneficial effect of statin. Therapy. independent of hyperlipidemia. To further clarify the differential benefit of lipid-dependent early statins, we sought to determine whether early statin therapy would be associated with a reduction in postoperative mortality or morbidity. In normolipidemic patients, the risk of death and MACE may increase after CABG.

Hyperlipidemia is a known risk factor for coronary artery disease and the progression of atherosclerosis, and an increase in some types of plasma fats has been shown to exacerbate atherosclerotic involvement of the coronary arteries [1]. There are different types of plasma fats, and among them, low-density lipoprotein (LDL) lipids have been directly and closely related to coronary artery disease, and high-density lipoproteins (HDL) are one of the strongest protective factors. Is the cause of atherosclerosis. Recent evidence suggests that a slight increase in glyceride levels increases the risk of coronary heart disease and accelerates coronary artery disease, as well as leading to new lesions and greater coronary artery involvement [2]. The association between serum lipid levels and coronary artery disease is well known, although studies on the role of lipoproteins as indicators of the severity of coronary artery involvement are still limited. The clinical manifestations of coronary artery disease have no independent and obvious relationship with the extent of coronary artery atherosclerosis, but the severity of the obstruction and the number of vessels involved are important in patients' treatment plans (figure 1).




Figure 1: Clinical Outcome in hyperlipidemia in CABG Candidate Patient

Finally, it is hypothesized that if one type of serum fat predicts the degree and severity of coronary artery involvement, it can be controlled by therapeutic strategies and the need for aggressive procedures such as catheterization and stenting or the need for

bypass surgery (Figure 2). The aim of this study was to evaluate the relationship between hyperlipidemia and serum levels of blood lipids and the number of coronary arteries involved in patients undergoing coronary artery bypass grafting.




Figure 2: Blocked Coronary Artery

Material and Methods

Study design: This is a cross-sectional descriptive study that was performed in 30 years in 2012-2015 in Shahid Madani Hospital (Tabriz University of Medical Sciences) with the participation of patients who are candidates for coronary artery bypass graft surgery. Sampling method was available in this study.

Inclusion and exclusion criteria: Inclusion criteria included consent to participate in the study, candidate for coronary artery bypass graft surgery and age over 18 years and exclusion criteria also included patients with a history of heart surgery, patients with congenital heart defects and death. Surgery was also one of the exclusion criteria of this study.

Methodology: By referring to the archives of the Open-Heart Surgery Department of Ghaem Hospital, the files of patients who underwent coronary artery bypass graft surgery in 2018 were reviewed and in addition to angiography, fat profile tests of patients were examined and then patients Based on the number of coronary arteries involved in angiography, they were divided into three groups: mono-vascular, bivascular and tricuspid. The prevalence of hyperlipidemia in these groups was compared to evaluate the relationship between hyperlipidemia and the number of coronary arteries involved.

Data analysis: After extracting the required information, the data were entered into SPSS software and data analysis and comparison in patient groups were performed using chi-square tests and analysis of variance.

Ethical Considerations: It is worth mentioning that the whole process of the present study was carried out after the approval and approval of the ethics committee of Tabriz University of Medical Sciences.

Results

In the study of 150 patients with coronary artery disease, candidates for CABG surgery, patients were divided into three groups based on the number of vessels involved in angiography, and it was observed that 4 patients (2.7%) had single involvement (Figure 3).




Figure 3: Number of lipid lowering

Vascular, 39 patients (26%) have 32 vascular involvement and 107 patients (71.3%) have vascular involvement. A total of 64 patients

(42.7%) were female and 86 patients (57.3%) were male and the mean age was 58.2 ± 10.86 years. Serum fat levels were assessed by the number of coronary arteries involved (Figure 4).




Figure 4: Mortality rate after hyperlipidemia candidate CABG

Discussion

Ischemic heart disease causes more death and disability than any other disease in developed and developing countries. These diseases have imposed economic costs on society and the individual, and with the urbanization of the developing world, the prevalence of the risk factor for IHD in these areas is increasing rapidly [5]. Widespread increases in IHD have been identified worldwide and are likely to become the most common cause of death worldwide by 2020. Also, the prevalence of this disease is increasing in our country [6]. It should be noted that several factors affect this disease, such as increased blood lipid levels, increased sugar and other risk factors, many of the risk factors that cause the disease can be prevented. For any intervention to reduce the risk factor and disease-causing factors, basic information is needed on the frequency of these risk factors in the community and patients, so this study was performed to determine the prevalence and type of hyperlipidemia in patients undergoing cardiac surgery [7]. In this study, 150 patients with coronary artery disease who were CABG candidates were divided into three groups based on the number of vessels involved in angiography: single-vessel, 2-vessel and 3-vessel involvement, and their serum fat levels were compared. Among 217 patients, single vascular involvement, 26% had 2 vascular involvement and 71.3% had 3 vascular involvement. In a study by Jin 1 and colleagues in China, 125 patients had much more single-vessel involvement, 113 had 2-vessel involvement, and 107 had 3-vessel involvement. In one study, among their patients, 33.4% had single-vessel involvement, 39.7% had two-vessel involvement and 27.4% had three-vessel involvement [8, 10-12]. Again, in the present study, the ratio of single-vessel involvement was much lower and CAD in our community It looks much wider and more

intense. In the present study, the overall male-to-female prevalence was 86 to 64, and like other studies in all groups, especially those with multiple vascular involvement, it was male. In the Ladia study in El Salvador, the male-to-female ratio was 62 to 38. In Penalva's study in 2008, 53.2% of her patients were male [9, 13-15]. The mean age of patients in this study was 58.2 years and the mean age was lower in the group of monovascular patients, although the differences were not significant. The mean age of CABG candidate patients was 56.3 years, which was similar to the present study [10, 16-18]. In all of our study groups, they were matched in terms of sex, age, smoking, and other MI. Although these risk factors were somewhat higher in multivariate conflict groups, the differences between the groups were not significant, so the distorting effect of these factors The danger was removed [11, 19-22]. In assessing blood lipid levels by considering the cut-off point of 130 for LDL for HLD 260 for HLD 260 for triglyceride and 200 for total cholesterol, in total patients increased LDL by 7.3%, decreased HDL by 10.7%, increased TG by 8% and increased cholesterol. Tom was reported to be 6.7% [8, 23-25]. Comparing the rate of hyperlipidemia in each case in different groups according to the number of vessels involved in angiography, it was observed that the prevalence of total cholesterol and LDL hyperlipidemia in multivascular patients is significantly higher than monovascular. In other studies, dyslipidemia, especially elevated cholesterol and LDL and decreased HDL, has been identified as a strong risk factor for coronary heart disease [26-28]. In one study, the ratio of total cholesterol to HDL was reported as the most important factor with the number of coronary arteries involved. In a study in El Salvador, the ratio of total cholesterol to HDL was significantly higher in patients with coronary artery disease [12, 29-31].

The main limitations of this study are its retrospective design and randomization. In addition, there were significant within-group differences in clinical and risk factors and within the study group itself. In addition, patients with hyperlipidemia who are not treated with statins are likely to have an incurable disease, rather than patients who have received "previously negative therapy" and are therefore at risk for other complications. Therefore, we explained these variables by carefully adjusting them using multiple risk-adjusted and quality-adjusted regression models. In addition, a small sample of statin-naïve patients over a follow-up period of more than 6 years suggests that statins are now widely available and highly effective. In addition, the effect of preoperative statin therapy (in terms of dose and duration) before CABG has not been evaluated and should be addressed in future studies.

Conclusion

Total cholesterol and LDL are important indicators for predicting the extent of coronary artery involvement. While the prevalence of elevated total cholesterol and elevated LDL are associated with multivascular involvement, the role of elevated LDL in multivessel coronary artery involvement is more effective than increased total cholesterol. Total cholesterol and LDL are among the predictors of the potential need for surgical intervention.

References

- [1] Mokhtari Ardekani AB, et al., miR-122 dysregulation is associated with type 2 diabetes mellitus-induced dyslipidemia and hyperglycemia independently of its rs17669 variant, *BioMed Research International*, **2022**, Article ID 5744008, [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [2] Mobaraki-Asl N, Ghavami Z, Gol MK. Development and validation of a cultural

competence questionnaire for health promotion of Iranian midwives. *Journal of education and health promotion*. **2019**;8:179.

- [3] Haghdoost M, Mousavi S, Gol MK, Montazer M. Frequency of Chlamydia trachomatis Infection in Spontaneous Abortion of Infertile Women During First Pregnancy Referred to Tabriz University of Medical Sciences by Nested PCR Method in 2015. *International Journal of Women's Health and Reproduction Sciences*. **2019**; 7(4): 526-30. [[Google Scholar](#)], [[Publisher](#)]
- [4] Gheisari R, Resalati F, Mahmoudi S, Golkari A, Mosaddad SA. Do Different Modes of Delivering Postoperative Instructions to Patients Help Reduce the Side Effects of Tooth Extraction? A Randomized Clinical Trial. *Journal of Oral and Maxillofacial Surgery*. **2018**;76(8):1652.e1-e7. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [5] Gheisari R, Doroodizadeh T, Estakhri F, Tadbir A, Soufdoost R, Mosaddad S. Association between blood groups and odontogenic lesions: a preliminary report. *Journal of Stomatology*. **2019**;72(6):269-73. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [6] Ghaibi E, et al., Comparison of Marital Satisfaction, Emotional Divorce and Religious Commitment among Nurses and Staff of Ahvaz Government Hospitals, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2022**, 1, 33-39. [[Google Scholar](#)], [[Publisher](#)],
- [7] Fathi A, Giti R, Farzin M. How different influential factors affect the color and translucency of y-ztp: a review of the literature. *Annals of Dental Specialty*. **2018**;1;6(3):338-41. [[Google Scholar](#)], [[Publisher](#)]
- [8] Eydi M, Golzari SE, Aghamohammadi D, Kolahdouzan K, Safari S, Ostadi Z. Postoperative management of shivering: A comparison of pethidine vs. ketamine. *Anesthesiology and Pain Medicine*; **2014**: 4(2),e15499 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [9] Elmi Sadr F, et al., The Risk for SARS-Cov-2 Virus Contamination through Surgical Smoking

- and Aerosolization by Laparoscopic Surgery: A Systematic Review, *Annals of the Romanian Society for Cell Biology* 25 (1), 6839 – 6852, [[Google Scholar](#)], [[Publisher](#)]
- [10] Eidy M, Ansari M, Hosseinzadeh H, Kolahdouzan K. Incidence of back pain following spinal anesthesia and its relationship to various factors in 176 patients. *Pakistan Journal of Medical Sciences*. **2010**; 26(4):778-781. [[Google Scholar](#)], [[Publisher](#)]
- [11] Eidi M, Kolahdouzan K, Hosseinzadeh H, Tabaqi R. A comparison of preoperative ondansetron and dexamethasone in the prevention of post-tympanoplasty nausea and vomiting. *Iranian Journal of Medical Sciences*. **2012**; 37(3):166-172. [[Google Scholar](#)], [[Publisher](#)]
- [12] Eghdam-Zamiri R, Gol MK. Effects of ginger capsule on treatment of nausea and vomiting in patients receiving cisplatin undergoing mastectomy: a randomized clinical trial. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. **2020**;22(11): 15-21.
- [13] Eghdam-Zamiri R, Gol MK. Effects of ginger capsule on treatment of nausea and vomiting in patients receiving cisplatin undergoing mastectomy: a randomized clinical trial. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. **2020**;22(11): 15-21. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [14] Ebadian B, Fathi A, Khodadad S. Comparison of the effect of four different abutment screw torques on screw loosening in single implant-supported prosthesis after the application of mechanical loading. *International Journal of Dentistry*. **2021**;19;2021: 3595064. [[Google Scholar](#)], [[Publisher](#)]
- [15] Dargahi R, Nazari B, Dorosti A, Charsouei S. Does coronavirus disease affect sleep disorders in the third trimester of pregnancy in women with low back pain? *International Journal of Women's Health and Reproduction Sciences*. **2021**; 9(4):268-273. [[Google Scholar](#)], [[Publisher](#)]
- [16] Darestani MN, Houshmand B, Mosaddad SA, Talebi M. Assessing the Surface Modifications of Contaminated Sandblasted and Acid-Etched Implants Through Diode Lasers of Different Wavelengths: An In-Vitro Study. *Photobiomodulation, Photomedicine, and Laser Surgery*. **2023**. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [17] Birman DH, Investigation of the Effects of Covid-19 on Different Organs of the Body, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2023**, 2 (1), 24-36 [[Google Scholar](#)], [[Publisher](#)]
- [18] Ansari Iari H, Jahed SH, Abbasi K, Alam M, Alihemmati M, Ghomi AJ, et al. In Vitro Comparison of the Effect of Three Types of Heat-Curing Acrylic Resins on the Amount of Formaldehyde and Monomer Release as well as Biocompatibility. *Advances in Materials Science and Engineering*. **2022**;2022:8621666. [[Google Scholar](#)], [[Publisher](#)]
- [19] Aghili SS, Pourzal A, Mosaddad SA, Amookhteh S. COVID-19 Risk Management in Dental Offices: A Review Article. *Open Access Maced J Med Sci*. **2022** Nov 04; 10(F):763-772. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20] Afshari A, Shahmohammadi R, Mosaddad SA, Pesteei O, Hajmohammadi E, Rahbar M, et al. Free-Hand versus Surgical Guide Implant Placement. *Advances in Materials Science and Engineering*. **2022**;2022:6491134. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21] Afshari A, Mosaddad SA, Alam M, Abbasi K, Darestani MN. Biomaterials and Biological Parameters for Fixed-Prosthetic Implant-Supported Restorations: A Review Study. *Advances in Materials Science and Engineering*. **2022**;2022:2638166. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [22] Abdollahi MH, Foruzan-Nia K, Behjati M, Bagheri B, Khanbabayi-Gol M, Dareshiri S and et al. The effect of preoperative intravenous paracetamol administration on postoperative

- fever in pediatrics cardiac surgery. Nigerian medical journal: journal of the Nigeria Medical Association. **2014**; 55(5): 379. [[Google Scholar](#)], [[Publisher](#)]
- [23] Abdollahi MH, Foruzan-Nia K, Behjati M, Bagheri B, Khanbabayi-Gol M, Dareshiri S and et al. The effect of preoperative intravenous paracetamol administration on postoperative fever in pediatrics cardiac surgery. Nigerian medical journal: journal of the Nigeria Medical Association. **2014**; 55(5): 379. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [24] Nazari B, Amani L, Ghaderi L, Gol MK. Effects of probiotics on prevalence of ventilator-associated pneumonia in multitrauma patients hospitalized in neurosurgical intensive care unit: a randomized clinical trial. Trauma Monthly. **2020**; 25(6): 262-268.
- [25] Susanabadi A, et al., A Systematic Short Review in Evaluate the Complications and Outcomes of Acute Severe of Pediatric Anesthesia, Journal of Chemical Reviews, **2021**, 3 (3), 219-231, [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26] Susanabadi A, et al., Evaluating the Outcome of Total Intravenous Anesthesia and Single Drug Pharmacological to Prevent Postoperative Vomiting: Systematic Review and Meta-Analysis, Annals of the Romanian Society for Cell Biology, **2021**, 25 (6), 2703-2716, [[Google Scholar](#)], [[Publisher](#)]
- [27] Golfeshan F, Ajami S, Khalvandi Y, Mosaddad SA, Nematollahi H. The Analysis of the Differences between the Influence of Herbal Mouthwashes and the Chlorhexidine Mouthwash on the Physical Characteristics of Orthodontic Acrylic Resin. Journal of Biological Research - Bollettino della Società Italiana di Biologia Sperimentale. **2020**;93(1). [[Google Scholar](#)], [[Publisher](#)]
- [28] Golfeshan F, Mosaddad SA, Babavalian H, Tebyanian H, Mehrjuyan E, Shakeri F. A Summary of Planarian Signaling Pathway for Regenerative Medicine. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences. **2022**;92(1):5-10. [[Google Scholar](#)], [[Publisher](#)]
- [29] Golfeshan F, Mosaddad SA, Ghaderi F. The Effect of Toothpastes Containing Natural Ingredients Such As Theobromine and Caffeine on Enamel Microhardness: An In Vitro Study. Evidence-Based Complementary and Alternative Medicine. **2021**;2021:3304543. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30] Danesh H, et al., Pharmacological Evaluation of Covid 19 Vaccine in Acute and Chronic Inflammatory Neuropathies, Journal of Medicinal and Chemical Sciences, **2022**, 561-570, [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [31] Kheradjoo H, et al., Mesenchymal stem/stromal (MSCs)-derived exosome inhibits retinoblastoma Y-79 cell line proliferation and induces their apoptosis, Molecular Biology Reports, **2023**, 50, 4217-4224, [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [32] Mahmoudi H, et al., Stem cell-derived nano-scale vesicles promotes the proliferation of retinal ganglion cells (RGCs) by activation PI3K/Akt and ERK pathway, Nanomedicine Research Journal, **2022**, 7(3), 288-293, [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [33] Haghdoost M, Mousavi S, Gol MK, Montazer M. Frequency of Chlamydia trachomatis Infection in Spontaneous Abortion of Infertile Women During First Pregnancy Referred to Tabriz University of Medical Sciences by Nested PCR Method in 2015. International Journal of Women's Health and Reproduction Sciences. **2019**; 7(4): 526-30. [[Google Scholar](#)], [[Publisher](#)]
- [34] Shirvani M, et al., The Diagnostic Value of Neutrophil to Lymphocyte Ratio as an Effective Biomarker for Eye Disorders: A Meta-Analysis, BioMed Research International, **2022**, Article ID 5744008, [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [35] Maalekipour M, Safari M, Berekatain M, Fathi A. Effect of adhesive resin as a modeling liquid on elution of resin composite restorations. *International Journal of Dentistry*. **2021**; 28;3178536. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [36] Mobaraki-Asl N, Ghavami Z, Gol MK. Development and validation of a cultural competence questionnaire for health promotion of Iranian midwives. *Journal of education and health promotion*. **2019**;8:179. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [37] Moharrami M, Anvari HM, Gheshlaghi LA, Nazari B. Preoperative education for pain relief after the lower limb joint replacement surgery: A systematic review and meta-analysis. *Trauma Monthly*; 26(1):52-60. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [38] Moharrami M, Nazari B, Anvari HM. Do the symptoms of carpal tunnel syndrome improve following the use of Kinesio tape? *Trauma Monthly*. **2021**; 26(4):228-234. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [39] Namanloo RA, Ommani M, Abbasi K, Alam M, Badkoobeh A, Rahbar M, et al. Biomaterials in Guided Bone and Tissue Regenerations: An Update. *Advances in Materials Science and Engineering*. **2022**;2022:2489399. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [40] Mosharraf R, Molaei P, Fathi A, Isler S. Investigating the Effect of Nonrigid Connectors on the Success of Tooth-and-Implant-Supported Fixed Partial Prostheses in Maxillary Anterior Region: A Finite Element Analysis (FEA). *International Journal of Dentistry*. **2021**; 12;2021: 5977994. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [41] Mosaddad, SA, Abdollahi Namanloo, R, Ghodsi, R, Salimi, Y, Taghva, M, Naeimi Darestani, M. Oral rehabilitation with dental implants in patients with systemic sclerosis: a systematic review. *Immun Inflamm Dis*. **2023**; 11:e812. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [42] Mosaddad SA, Yazdanian M, Tebyanian H, Tahmasebi E, Yazdanian A, Seifalian A, et al. Fabrication and properties of developed collagen/strontium-doped Bioglass scaffolds for bone tissue engineering. *Journal of Materials Research and Technology*. **2020**;9(6):14799-817. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [43] Mosaddad SA, Salari Y, Amookhteh S, Soufdoost RS, Seifalian A, Bonakdar S, et al. Response to Mechanical Cues by Interplay of YAP/TAZ Transcription Factors and Key Mechanical Checkpoints of the Cell: A Comprehensive Review. *Cell Physiol Biochem*. **2021**;55(1):33-60. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [44] Mosaddad SA, Rasoolzade B, Namanloo RA, Azarpira N, Dortaj H. Stem cells and common biomaterials in dentistry: a review study. *Journal of Materials Science: Materials in Medicine*. **2022**;33(7):55. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [45] Mosaddad SA, Namanloo RA, Aghili SS, Maskani P, Alam M, Abbasi K, et al. Photodynamic therapy in oral cancer: a review of clinical studies. *Medical Oncology*. **2023**;40(3):91. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [46] Mosaddad SA, Gheisari R, Erfani M. Oral and maxillofacial trauma in motorcyclists in an Iranian subpopulation. *Dental Traumatology*. **2018**;34(5):347-52. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [47] Mosaddad SA, Beigi K, Doroodizadeh T, Haghnegahdar M, Golfeshan F, Ranjbar R, et al. Therapeutic applications of herbal/synthetic/bio-drug in oral cancer: An update. *Eur J Pharmacol*. **2021**;890:173657. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [48] Nazari B, Amani L, Ghaderi L, Gol MK. Effects of probiotics on prevalence of ventilator-associated pneumonia in multitrauma patients hospitalized in neurosurgical intensive care unit: a randomized clinical trial. *Trauma*

- Monthly. **2020**; 25(6): 262-268. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [49] Birmangi S, A Review of the Effect of Corona on the Human Brain – Short Review, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2022**, 1 (3), 80-87 [[Google Scholar](#)], [[Publisher](#)]
- [50] Margy S, A Review of the Effect of Brain imaging- Short Review, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2022**, 1 (3), 88-99 [[Google Scholar](#)], [[Publisher](#)]
- [51] Musaei S, The Effect of Pregnancy on the Skin, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2023**, 2(1), 17-23. [[Google Scholar](#)], [[Publisher](#)]
- [52] Nazardani SZ, et al., A comprehensive evaluation of the Sports Physiotherapy curriculum, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2023**, 2, 10-16. [[Google Scholar](#)], [[Publisher](#)]
- [53] Yazdani M, Rahmani A, Tahmasebi E, Tebyanian H, Yazdani A, Mosaddad SA. Current and Advanced Nanomaterials in Dentistry as Regeneration Agents: An Update. *Mini Reviews in Medicinal Chemistry*. **2021**;21(7):899-918. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [54] Tosan F, Rahnama N, Sakhaei D, Fathi AH, Yari A. Effects of doping metal nanoparticles in hydroxyapatite in Improving the physical and chemical properties of dental implants. *Nanomedicine Research Journal*. **2021**;1;6(4):327-36. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [55] Tahmasebi E, Alam M, Yazdani M, Tebyanian H, Yazdani A, Seifalian A, et al. Current biocompatible materials in oral regeneration: a comprehensive overview of composite materials. *Journal of Materials Research and Technology*. **2020**;9(5):11731-55. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [56] Sarejloo SH, et al., Neutrophil-to-Lymphocyte Ratio and Early Neurological Deterioration in Stroke Patients: A Systematic Review and Meta-Analysis, **2022**, Article ID 8656864 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [57] Saliminasab M, Jabbari H, Farahmand H, Asadi M, Soleimani M, Fathi A. Study of antibacterial performance of synthesized silver nanoparticles on Streptococcus mutans bacteria. *Nanomedicine Research Journal*. 2022 Oct 1;7(4):391-6. [[Google Scholar](#)], [[Publisher](#)]
- [58] Mahkooyeh SA, et al., Chemical laboratory findings in children with covid-19: A systematic review and meta-analysis, *Eurasian Chemical Communications*, **2022**, 338-346, [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [59] Ahmadi SE, et al., Succinct review on biological and clinical aspects of Coronavirus disease 2019 (COVID-19), *Romanian Journal of Military Medicine*, **2022**,356-365, [[Google Scholar](#)], [[Publisher](#)]

This journal is a double-blind peer-reviewed journal covering all areas in Chemistry, Medicinal and Petroleum. EJCMPR is published quarterly (6 issues per year) online and in print. Copyright © 2022 by ASC ([Amir Samimi Company](#)) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.