

Original Article: Predicts factor in Polycystic Ovary Syndrome: An Evidence-based Study and Invitation from the Chief Editor

Fakhreddin Rasouli

Medical Doctor, Medicine Faculty, Urmia University of Medical Science, Urmia, Iran (OCID: 0000-0002-3060-5065)

Use your device to scan and read the article online



Citation F Rasouli, Predicts factor in Polycystic Ovary Syndrome: An evidence-based study and invitation from the chief editor, *EJCMPR*. 2024; 3(2):644-650



<https://doi.org/EJCMPR/20240713>

Article info:

Received: 27 February 2024

Accepted: 06 Jun 2024

Available Online:

ID: EJCMPR-2406-1179

Checked for Plagiarism: Yes

Peer Reviewers Approved by:

Dr. Frank Rebout

Editor who Approved Publication:

Dr. Frank Rebout

Keywords:

Polycystic Ovary Syndrome, risk factor, evidence-based study

ABSTRACT

Polycystic Ovary Syndrome (PCOS) is one of the most common endocrine disorders affecting women of reproductive age, with a prevalence of approximately 6-10% worldwide. Characterized by a range of symptoms including irregular menstrual cycles, hyperandrogenism, and polycystic ovaries, PCOS presents a complex challenge for both diagnosis and management. The syndrome's etiology is multifactorial, involving genetic, environmental, and lifestyle factors. This comprehensive review aims to explore the various predictive factors associated with PCOS, encompassing genetic predispositions, hormonal imbalances, metabolic dysfunctions, lifestyle influences, and potential environmental triggers. Characterized by symptoms such as menstrual irregularities, hyperandrogenism, and polycystic ovaries, PCOS presents a broad spectrum of clinical manifestations. The multifaceted nature of PCOS makes it a challenge to diagnose and manage effectively. Understanding the predictive factors involved in PCOS is crucial for early identification and personalized treatment.

Introduction

Polycystic Ovary Syndrome (PCOS) is a heterogeneous endocrine disorder that affects a significant percentage of women in their reproductive years [1]. Characterized by symptoms such as menstrual irregularities, hyperandrogenism, and polycystic ovaries [2], PCOS presents a broad spectrum of clinical manifestations. The multifaceted nature of PCOS makes it a challenge

to diagnose and manage effectively. Understanding the predictive factors involved in PCOS is crucial for early identification and personalized treatment. This extensive review examines the genetic predispositions, hormonal imbalances, metabolic dysfunctions, lifestyle influences, and environmental triggers that contribute to the development and progression of PCOS [3].

*Corresponding Author: Fakhreddin Rasouli (Email: RFakhreddin90@gmail.com)

Genetic Predispositions

Genetic factors play a pivotal role in the etiology of PCOS. Family studies have demonstrated a higher prevalence of PCOS among first-degree relatives, suggesting a strong hereditary component. The heritability of PCOS is estimated to be between 60% and 80%, indicating a significant genetic influence. Several genes have been implicated in the pathogenesis of PCOS, including those involved in insulin signaling, androgen biosynthesis, and folliculogenesis [4]. Genome-wide association studies (GWAS) have identified multiple genetic loci associated with PCOS [5].

Notable genes include LHCGR, which encodes the luteinizing hormone/choriogonadotropin receptor, and FSHR, which encodes the follicle-stimulating hormone receptor. Variants in these genes may disrupt normal ovarian function, leading to the hormonal imbalances observed in PCOS. Another critical gene is INSR, encoding the insulin receptor, which is linked to the insulin resistance commonly seen in PCOS patients [6]. Studies have also highlighted the role of epigenetic modifications in PCOS. Epigenetic changes, such as DNA methylation and histone modification, can alter gene expression without changing the underlying DNA sequence. These modifications can be influenced by environmental factors and may contribute to the heritability and phenotypic variability of PCOS.

Hormonal Imbalances

Hormonal dysregulation is a central feature of PCOS, with hyperandrogenism being one of the most consistent abnormalities. Elevated androgen levels, including testosterone, androstenedione, and dehydroepiandrosterone sulfate (DHEAS), are commonly observed in women with PCOS. Hyperandrogenism manifests clinically as hirsutism, acne, and alopecia, contributing significantly to the physical and psychological burden of the disorder [7-9].

The source of excess androgens in PCOS is multifactorial. Increased ovarian androgen production is a primary contributor, driven by the hypersecretion of luteinizing hormone (LH). In PCOS, there is an abnormal LH to follicle-stimulating hormone (FSH) ratio, with elevated LH levels stimulating the theca cells of the ovaries to produce more androgens. Additionally [10-12], insulin resistance and hyperinsulinemia exacerbate hyperandrogenism by increasing ovarian androgen synthesis and reducing hepatic production of sex hormone-binding globulin (SHBG), which binds androgens and decreases their bioavailability [13].

Another hormonal imbalance observed in PCOS is the dysfunction of the hypothalamic-pituitary-ovarian (HPO) axis. The dysregulation of this axis results in disrupted folliculogenesis and anovulation, leading to menstrual irregularities. Elevated LH levels, coupled with insufficient FSH secretion, impede the normal maturation of ovarian follicles, contributing to the formation of cystic ovaries.

Metabolic Dysfunctions

Metabolic abnormalities are prevalent in women with PCOS, with insulin resistance being a key feature. Insulin resistance occurs when cells in the body become less responsive to the action of insulin, resulting in elevated blood glucose levels and compensatory hyperinsulinemia. Insulin resistance is present in approximately 70% of women with PCOS, irrespective of body mass index (BMI).

Hyperinsulinemia exacerbates hyperandrogenism by promoting ovarian androgen production and inhibiting SHBG synthesis. This metabolic dysfunction creates a vicious cycle that perpetuates the clinical manifestations of PCOS. Moreover, insulin resistance increases the risk of developing type 2 diabetes mellitus, cardiovascular disease, and metabolic syndrome in women with PCOS.

These comorbidities highlight the importance of addressing metabolic dysfunctions in the management of PCOS.

Adipose tissue dysfunction is also a significant contributor to the metabolic disturbances in PCOS. Adipose tissue, particularly visceral fat, is metabolically active and secretes various adipokines and inflammatory cytokines. In women with PCOS, adipose tissue dysfunction leads to increased levels of pro-inflammatory cytokines, such as tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6), which further exacerbate insulin resistance and contribute to chronic low-grade inflammation.

Lifestyle Influences

Lifestyle factors, including diet, physical activity, and stress, play a crucial role in the development and progression of PCOS. Obesity, particularly central obesity, is commonly associated with PCOS and exacerbates its clinical features. Excess adipose tissue contributes to insulin resistance and hyperandrogenism, making weight management a critical component of PCOS treatment [14].

Dietary patterns high in refined carbohydrates, saturated fats, and low in fiber are linked to increased insulin resistance and inflammation, which can worsen PCOS symptoms. Conversely, a balanced diet rich in whole grains, lean proteins, healthy fats, and vegetables can improve insulin sensitivity and reduce androgen levels. Nutritional interventions, such as the Mediterranean diet and low glycemic index (GI) diets, have shown promise in managing PCOS symptoms and improving metabolic parameters [16].

Regular physical activity is essential for improving insulin sensitivity, promoting weight loss, and reducing androgen levels in women with PCOS. Exercise not only aids in weight management but also enhances overall metabolic health. Both aerobic and resistance training have been shown to be beneficial in

improving the metabolic and reproductive outcomes in PCOS.

Stress management is another critical aspect of lifestyle intervention in PCOS. Chronic stress can disrupt the hypothalamic-pituitary-adrenal (HPA) axis, leading to increased cortisol levels and exacerbating hormonal imbalances. Stress reduction techniques, such as mindfulness, yoga, and cognitive-behavioral therapy (CBT), can help mitigate the impact of stress on PCOS and improve overall well-being.

Environmental Triggers

Environmental factors, particularly exposure to endocrine-disrupting chemicals (EDCs), have been implicated in the pathogenesis of PCOS. EDCs are exogenous compounds that interfere with hormone signaling pathways, potentially leading to reproductive disorders. Common EDCs include bisphenol A (BPA), phthalates, and pesticides [15].

Studies have shown higher levels of BPA in women with PCOS compared to controls. BPA is known to mimic estrogen and disrupt normal endocrine function, which may contribute to the hormonal imbalances observed in PCOS. Animal studies have demonstrated that prenatal and early-life exposure to BPA can result in PCOS-like phenotypes, suggesting that EDC exposure may have long-term effects on reproductive health.

Phthalates, commonly found in plastics and personal care products, have also been linked to PCOS. These chemicals can disrupt androgen and estrogen signaling, potentially exacerbating the hyperandrogenism and menstrual irregularities seen in PCOS. The exact mechanisms by which EDCs influence PCOS development remain to be fully elucidated, but they likely involve complex interactions with genetic and environmental factors.

Psychological and Emotional Factors

PCOS has a profound impact on psychological and emotional well-being. Women with PCOS are at an increased risk of experiencing mood disorders, including depression and anxiety. The chronic nature of PCOS, along with its physical manifestations such as hirsutism, acne, and obesity, can lead to significant body image issues and reduced self-esteem [17].

The hormonal imbalances associated with PCOS, particularly elevated androgen levels, may directly contribute to mood disturbances. Additionally, the metabolic complications of PCOS, including insulin resistance and obesity, are linked to increased inflammation, which has been associated with depression.

Managing the psychological aspects of PCOS is essential for improving the overall quality of life. Integrative approaches that address both the physical and emotional aspects of PCOS, such as cognitive-behavioral therapy (CBT), support groups, and lifestyle modifications, can help mitigate the psychological burden of the disorder [18].

Reproductive Implications

PCOS is a leading cause of infertility due to chronic anovulation. The hormonal imbalances, particularly the elevated LH to FSH ratio, disrupt normal folliculogenesis and ovulation. Women with PCOS often present with irregular menstrual cycles, making it challenging to predict ovulation and conceive naturally.

Ovulation induction therapies, such as clomiphene citrate and letrozole, are commonly used to stimulate ovulation in women with PCOS. In cases where these treatments are ineffective, assisted reproductive technologies (ART) such as in vitro fertilization (IVF) may be necessary. Despite the challenges, many women with PCOS can achieve successful pregnancies with appropriate medical intervention.

Pregnancy in women with PCOS is associated with an increased risk of complications,

including gestational diabetes, preeclampsia, and preterm birth. Therefore, close monitoring and management of metabolic and hormonal parameters during pregnancy are essential to ensure maternal and fetal health [19-21].

Innovative Therapeutic Approaches

Emerging therapies and novel approaches are being explored to manage PCOS more effectively. Inositol supplements, particularly myo-inositol and D-chiro-inositol, have shown promise in improving insulin sensitivity and reducing androgen levels [22-24]. These supplements mimic the action of insulin and enhance its signaling pathways, offering a potential alternative or adjunct to traditional treatments.

Anti-androgen medications, such as spironolactone and flutamide, are used to manage hyperandrogenic symptoms like hirsutism and acne. These medications block androgen receptors and reduce the effect of excess androgens on the skin and hair follicles. Metformin, a well-known insulin sensitizer, is commonly prescribed to women with PCOS to improve insulin resistance and restore menstrual regularity. It also has beneficial effects on weight management and metabolic health [25].

Bariatric surgery is considered for women with PCOS who are morbidly obese and have not responded to conventional weight loss strategies. Bariatric surgery can lead to significant weight loss, improved insulin sensitivity, and normalization of menstrual cycles, thereby ameliorating the symptoms of PCOS.

Future Directions in PCOS Research

Ongoing research aims to further elucidate the underlying mechanisms of PCOS and develop more effective treatments. Advances in genetic and epigenetic research hold promise for

identifying new therapeutic targets and understanding the hereditary aspects of PCOS.

The role of the gut microbiome in PCOS is an emerging area of interest [26]. Dysbiosis, or an imbalance in the gut microbiota, has been linked to metabolic disorders, including insulin resistance and obesity. Studies suggest that the gut microbiome may influence the hormonal and metabolic features of PCOS, offering a potential avenue for novel treatments such as probiotics and dietary interventions.

Precision medicine, which tailors treatment based on individual genetic, environmental, and lifestyle factors, is an exciting frontier in PCOS management. By understanding the unique profile of each patient, healthcare providers can offer more personalized and effective treatment strategies [27].

Conclusion

Polycystic Ovary Syndrome is a complex disorder with a multifactorial etiology involving genetic, hormonal, metabolic, lifestyle, and environmental factors. Understanding the predictive factors associated with PCOS is essential for early diagnosis, personalized treatment, and effective management. Comprehensive approaches that address the physical, psychological, and emotional aspects of PCOS are crucial for improving the quality of life for women affected by this disorder. Continued research and innovative therapeutic strategies hold promise for better understanding and managing PCOS in the future.

References

- [1]Jiang S, Zhang Y, Alsaikhan F, Jalil AT, Gol MK, Tarighatnia A. A meta-analysis review of the effect of Zn-doped synthetic polymer materials on bone regeneration. *Journal of Drug Delivery Science and Technology*. 2022 Oct 1;76: 103792 [Google Scholar], [Publisher], [Crossref]
- [2]Gol MK, Payami S, Lotfi A. Study of the effect of ear acupressure on stress and serum cortisol

level before rhinoplasty surgery: a randomized clinical trial. *Crescent Journal of Medical & Biological Sciences*. 2020 Apr 1;7(2). [Google Scholar], [Publisher]

[3]Khanbabayi Gol M, Arefi N, Jafari M, Farzin H, Aghamohammadi D. Prevalence of Post-Related Infections and Their Predisposing Factors in Women with Breast Cancer under Chemotherapy. *Iranian Quarterly Journal of Breast Disease*. 2018 Sep 10;11(2):7-15. [Google Scholar], [Publisher], [Crossref]

[4]Piroozkhah, Moein, et al. "Guanylate cyclase-C Signaling Axis as a theragnostic target in colorectal cancer: a systematic review of literature." *Frontiers in Oncology* 13 (2023). [Google Scholar], [Publisher], [Crossref]

[5]AZHOUGH, RAMIN, et al. "Surgical Outcome and Complications of Lateral Internal Sphincterotomy in Chronic Anal Fissure-A Cross-sectional Study." *Journal of Clinical & Diagnostic Research* 15.12 (2021). [Google Scholar], [Publisher], [Crossref]

[6]Jalali, Pooya, et al. "UBE2C: A pan-cancer diagnostic and prognostic biomarker revealed through bioinformatics analysis." *Cancer Reports* 7.4 (2024): e2032. [Google Scholar], [Publisher], [Crossref]

[7]Azough, Ramin, et al. "Pilonidal sinus surgery with mechanical drilling technique." *Indian Journal of Surgery* 82 (2020): 824-827. [Google Scholar], [Publisher], [Crossref]

[8]Azough, Ramin, et al. "Intradermal methylene blue analgesic application in post hemorrhoidectomy pain management: a randomized controlled trial." *Frontiers in Surgery* 11 (2024): 1354328. [Google Scholar], [Publisher], [Crossref]

[9]K Hadavand Mirzaei, et al., Application of Robotic Instruments in Hip Arthroplasty Surgery Based on Practical Tips a Systematic Review, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, 2024, 3 (2), 593-609 [Google Scholar], [Publisher], [Crossref]

- [10] S Sourili, What are the radiological criteria for identifying the Corona virus in CT scan images?, Eurasian Journal of Chemical, Medicinal and Petroleum Research (EJCMPR), **2024**, 3(2), 577-593 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [11] S Soubkari, Covid CT Chest Finding, Eurasian Journal of Chemical, Medicinal and Petroleum Research (EJCMPR), **2024**, 3(2), 563-576 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [12] S Asadimehr, Examining Drilling Problems and Practical Solutions Regarding them, Eurasian Journal of Chemical, Medicinal and Petroleum Research, **2024**, 3 (2), 552-562 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [13] S Asadimehr, Investigating the Use of Drilling Mud and the Reasons for its Use, Eurasian Journal of Chemical, Medicinal and Petroleum Research, **2024**, 3 (2), 543-551 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [14] S Soubhak, Nursing actions in patients undergoing laryngectomy in relation to reducing anxiety and depression, Eurasian Journal of Chemical, Medicinal and Petroleum Research, **2024**, 3 (2), 530-542 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [15] S Soubhak, Nursing actions in patients undergoing laryngectomy in relation to reducing anxiety and depression, Eurasian Journal of Chemical, Medicinal and Petroleum Research, **2024**, 3 (2), 530-542
- [16] R.F Alaouadi, Frequency of V. cholera in Babylon Province. Journal of Babylon University/Pure and Applied Sciences, **2014**, 22, 2579-2590 [[Google Scholar](#)], [[Publisher](#)]
- [17] WF Tawfeeq, et al., Detection of hepatitis C virus infection and genotypes among seropositive blood donors by polymerase chain reaction in babylon Governorate, Iraq. Medical Journal of Babylon. **2013** 10(1):25-37 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [18] RF Al-Jobouri., Assessment of immune response in patients with type 2 Diabetes mellitus. Ann Trop Med Public Health. **2020** 23(2):51-7 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [19] R.F. Alaouadi., Comparative phenotypic and genotypic study of MSSA and MRSA wound infections in Babylon, Al-Qadisiyah. J Vet Med Sci., **2015** 14, 23-33 [[Google Scholar](#)], [[Publisher](#)]
- [20] RF Alaouadi., Effect of age and sex on immunological activation in relation to infection with Echinococcus granulosus. J University Babylon. **2016** 24: 1935-46 [[Google Scholar](#)], [[Publisher](#)]
- [21] RF Aboqader et al., INCREASE IL-17 α AND REDUCED IFN- γ /IL-4 RATIO ARE NOVEL RESPONSIBLE FACTORS FOR PERSISTENCE OF ENTAMOEBA HISTOLYTICA INFECTION. Biochemical & Cellular Archives, **2017** 19 (2), 817-823 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [22] SA Alkarady., et al., Assessment the Level of Interleukine-5 and Eosinophils Count in Allergic Rhinitis and Asthma Patients Journal of Babylon University/Pure and Applied Sciences, **2016** 5(24): 1221-1226 [[Google Scholar](#)], [[Publisher](#)]
- [23] RF AL-Aouadi., et al., Study of Immunological Response to Early and Late Stages of Hydatid Cyst Formation, Journal of University of Babylon, 2016 24(5) 1433-1446 [[Google Scholar](#)], [[Publisher](#)]
- [24] RF Al-Awadi., et al., Assessment of Immunopathological Response of Serum Immunoglobulins and Protective Role of Complement Components in Iraqi Patients with Hydatidosis. Iraqi postgraduate medical journal, **2008** 7(4)322-326 [[Google Scholar](#)], [[Publisher](#)]
- [25] RF Alaouadi, BK Salman Saqban., AKS Gene Expression of Formyl Peptide Receptor 1 in Eosinophils of Iraqi Patients with Extrinsic Allergic Asthma, 2014 13(3): 08-12 [[Google Scholar](#)], [[Publisher](#)]
- [26] FH Al Baw., RF Aboqader., Interplay of IL-2 and IL-17 in immunopathogenesis of vitiligo

Middle Euphrate, Iraq, 2019 22(12):227-230
[[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
[27] RF Aboqader., MH Al-Hasnawy., IL-8 is a
valuable laboratory marker for detection of

amoebiasis severity, **2019** Corpus ID:
210961635 [[Google Scholar](#)], [[Publisher](#)],
[[Crossref](#)]

This journal is a double-blind peer-reviewed journal covering all areas in Chemistry, Medicinal and Petroleum. EJCMR is published quarterly (6 issues per year) online and in print. Copyright © **2024** by ASC ([Amir Samimi Company](#)) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.