

Original Article: Abdominoplasty Surgery



Mehrnoosh Rassam^{1*}, Bahram Davoudi²

¹General Surgeon residence, Department of General Surgery, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran (ORCID: 0000-0009-5237-1410)

²Medical Doctor, Islamic Azad University Tehran Medical Sciences, Tehran, Iran (ORCID: 0009-0004-3141-2937)

Use your device to scan and read the article online



Citation M Rassam, B Davoudi, **Abdominoplasty Surgery**, *EJCMPR*. 2024; 3(3):703-713



<https://doi.org/EJCMPR/20240116>

Article info:

Received: 27 February 2024

Accepted: 07 Jun 2024

Available Online:

ID: EJCMPR-2406-1183

Checked for Plagiarism: Yes

Peer Reviewers Approved by:

Dr. Frank Rebout

Editor who Approved Publication:

Dr. Frank Rebout

Keywords:

Abdominoplasty, Postoperative Care, Tummy Tuck

ABSTRACT

Abdominoplasty, commonly known as a "tummy tuck," is a widely sought-after cosmetic surgery aimed at enhancing the abdominal contour by removing excess skin and fat, and tightening the underlying muscles. The evolution of abdominoplasty techniques has led to a variety of approaches tailored to meet diverse patient needs and goals. Key techniques include the full abdominoplasty, which addresses significant skin and muscle laxity; the mini abdominoplasty, suited for patients with minimal excess skin below the navel; the extended abdominoplasty, which also targets the flanks and lower back; and the circumferential abdominoplasty, ideal for massive weight loss patients requiring comprehensive body contouring. Combining liposuction with abdominoplasty has become popular for refining results and achieving a more sculpted appearance. Advances such as progressive tension sutures and laser-assisted liposuction have further improved outcomes by minimizing complications and enhancing skin tightening. The selection of technique is based on a thorough preoperative assessment, ensuring personalized surgical plans that align with patient expectations. Effective postoperative care, including pain management and supportive garments, is crucial for optimal recovery and results. As abdominoplasty techniques continue to advance, they offer significant improvements in both aesthetic appearance and patient satisfaction, contributing to enhanced overall well-being.

Introduction

Abdominoplasty, commonly known as a "tummy tuck," is a surgical procedure designed to remove excess fat and skin from the abdominal area and to tighten the muscles [1-3], resulting in a firmer and more aesthetically pleasing abdominal profile [4-6]. This cosmetic surgery has become increasingly popular due to its dual benefits of enhancing

physical appearance and improving the functionality of the abdominal region. The advancements in surgical techniques over the years have significantly improved the safety and efficacy of abdominoplasty [7-9], making it a viable option for many individuals seeking body contouring solutions [10-12].

The history of abdominoplasty dates back to the early 20th century, initially focused on medical issues like hernia repair [13-15]. As surgical

*Corresponding Author: Mehrnoosh Rassam (Email: Rassam_mhrnsh@gmail.com)

techniques progressed, the procedure evolved to address aesthetic concerns. By the mid-20th century, abdominoplasty had begun to gain recognition as a cosmetic surgery aimed at enhancing the abdominal appearance. Today, it is a well-established procedure, thanks to the refinement of techniques and a deeper understanding of human anatomy, providing reliable and satisfactory results for patients [16]. Abdominoplasty is performed for a variety of reasons, which can be categorized into aesthetic and medical indications. Aesthetic reasons often include post-pregnancy changes, significant weight fluctuations, and the effects of aging. Pregnancy can cause significant stretching of the abdominal muscles and skin, often resulting in a protruding abdomen and stretch marks that do not revert to their pre-pregnancy state [17-19]. Substantial weight loss can leave behind loose, sagging skin that fails to retract, despite maintaining a stable weight. As people age, the skin loses its elasticity and muscles may weaken, leading to a less toned abdominal area. Medical indications for abdominoplasty can involve conditions like rectus diastasis, hernias, and the need for functional improvement [20-22]. Rectus diastasis, a condition involving the separation of the abdominal muscles, often occurs due to pregnancy or significant weight fluctuations, leading to a bulging abdomen that may benefit from surgical correction. Some patients with abdominal hernias can have these repaired simultaneously during an abdominoplasty. Additionally, excess abdominal skin can cause issues such as skin irritation, infections, and difficulty with physical activities, which can be alleviated through abdominoplasty [23-25].

There are several types of abdominoplasty procedures designed to address different patient needs and desired outcomes. The most common form is the full abdominoplasty, which involves a horizontal incision between the pubic hairline and the navel. Excess skin and fat are

removed, abdominal muscles are tightened, and the navel is repositioned.

A mini abdominoplasty, suitable for individuals with less extensive excess skin and fat localized below the navel, involves a shorter incision and typically does not reposition the navel [26].

An extended abdominoplasty addresses excess skin and fat in the abdominal area as well as the flanks, with a longer incision extending around the hips for comprehensive contouring. A circumferential abdominoplasty, often performed on patients who have experienced massive weight loss, involves an incision around the entire waistline, addressing excess skin and fat on the abdomen, back, and hips [27-29].

Before undergoing abdominoplasty, patients undergo a thorough medical evaluation to ensure they are suitable candidates. This includes reviewing medical history, conducting physical examinations, and discussing goals and expectations. Preoperative instructions typically involve lifestyle modifications, such as quitting smoking, avoiding certain medications, and maintaining a stable weight.

On the day of surgery, the patient is placed under general anesthesia for comfort and safety. The surgeon makes the necessary incisions based on the type of abdominoplasty. For a full abdominoplasty, a horizontal incision is made between the pubic hairline and the navel. The skin is then separated from the abdominal wall to reveal the underlying muscles [30].

These muscles are pulled together and stitched into a new position, creating a firmer abdominal wall and a narrower waistline. Excess skin is removed, and the remaining skin is pulled down and sutured together [31].

A new opening for the navel is created if needed, and the navel is brought out through this new incision and sutured in place. For mini or extended abdominoplasties, the incision length and muscle repair extent vary, but the basic principles remain the same. In some cases,

liposuction may be performed in conjunction with abdominoplasty to enhance contouring.

Recovery from abdominoplasty varies depending on the extent of the surgery and individual patient factors. Generally, patients can expect to stay in the hospital for a few hours to a day after the procedure, depending on their surgeon's recommendations [32].

During the initial recovery phase, patients typically experience pain and discomfort, which can be managed with prescribed pain medications. Bruising and swelling are also common and will gradually subside over time. Patients are encouraged to start walking as soon as possible to promote blood circulation and reduce the risk of blood clots. However, strenuous activities should be avoided for several weeks. Proper wound care is crucial to prevent infection and ensure optimal healing. Patients are instructed on how to care for their incisions and drains if they are placed [33].

While initial swelling may subside within a few weeks, residual swelling can persist for several months. The final results of abdominoplasty are typically visible around six months to a year post-surgery. Scars from abdominoplasty are permanent but will fade over time. Surgeons make incisions in locations that can be easily concealed by underwear or swimsuits. Maintaining a healthy lifestyle, including a balanced diet and regular exercise, is essential to preserve the results of the surgery.

As with any surgical procedure, abdominoplasty carries potential risks and complications. It is important for patients to be aware of these and discuss them thoroughly with their surgeon before deciding to undergo the procedure. Common risks include infection, excessive bleeding, scarring, seroma (fluid accumulation under the skin), and numbness due to nerve damage during surgery. More severe complications, though rare, include deep vein thrombosis (DVT) and pulmonary embolism

(PE), which involve blood clots that can travel to the lungs.

Poor wound healing can also occur, especially in patients who smoke or have diabetes, leading to noticeable scars or the need for revision surgery. In very rare cases, the blood supply to the skin can be compromised, leading to tissue death and necessitating further surgical intervention [34]. The decision to undergo abdominoplasty is deeply personal and can have significant psychological and emotional implications. For many individuals, the procedure offers a substantial boost in self-esteem and body image, leading to an enhanced quality of life. However, it is important for patients to have realistic expectations and understand the limitations of the surgery.

Psychological readiness and a clear understanding of the potential risks and benefits are crucial for achieving a satisfying outcome. In conclusion, abdominoplasty is a transformative surgical procedure that addresses both aesthetic and functional concerns of the abdominal area. With careful consideration, thorough preparation, and adherence to postoperative care instructions, patients can achieve a smoother, firmer [35], and more aesthetically pleasing abdominal profile, significantly improving their overall well-being.

Anesthesiology in abdominoplasty surgery

Anesthesiology plays a pivotal role in abdominoplasty surgery, ensuring patient comfort, safety, and optimal surgical outcomes. Abdominoplasty, commonly known as a "tummy tuck," is a major surgical procedure aimed at removing excess skin and fat from the abdominal area and tightening the underlying muscles to create a smoother, firmer abdominal profile.

This complex surgery, which involves significant tissue manipulation and potential blood loss, requires meticulous anesthetic management to

mitigate risks, manage pain, and enhance the overall surgical experience [36].

Understanding the intricacies of anesthesiology in abdominoplasty is crucial for both patients and healthcare providers to appreciate the comprehensive care involved in this transformative procedure.

Anesthesiology is a specialized field of medicine focused on the use of anesthesia and perioperative care, encompassing preoperative assessment, intraoperative management, and postoperative recovery [37].

The anesthesiologist, a physician specialized in this field, is responsible for administering anesthetics, monitoring the patient's vital signs, and ensuring a smooth transition through the various stages of surgery. In abdominoplasty, the role of the anesthesiologist extends beyond merely "putting the patient to sleep"; it involves a thorough understanding of the patient's medical history, surgical plan, and potential complications that could arise during and after the procedure.

The process begins with a detailed preoperative assessment, where the anesthesiologist evaluates the patient's overall health, medical history, and specific factors that could influence anesthesia administration. This assessment includes reviewing any chronic conditions, previous surgical experiences, allergies, medications, and lifestyle factors such as smoking or alcohol use. Patients are also typically advised to avoid certain medications and follow specific guidelines to prepare for anesthesia [38].

This comprehensive evaluation helps in formulating an individualized anesthetic plan, tailored to the patient's unique needs and the specifics of the abdominoplasty procedure.

There are various types of anesthesia that can be utilized in abdominoplasty, each with its advantages and considerations. General anesthesia is the most commonly employed method, rendering the patient completely

unconscious and insensitive to pain. This approach is preferred for its ability to provide complete immobility and control over the patient's airway and breathing, essential for a lengthy and invasive procedure like abdominoplasty [39]. The administration of general anesthesia involves the use of intravenous drugs and inhaled gases to induce and maintain unconsciousness, alongside the insertion of a breathing tube to secure the airway.

Regional anesthesia, such as epidural or spinal anesthesia, may also be considered in some cases. These methods involve the injection of anesthetic agents near the spinal cord, numbing the lower part of the body while the patient remains awake or sedated. Although less common for abdominoplasty, regional anesthesia can offer benefits such as reduced postoperative pain and quicker recovery of consciousness [40]. However, the choice between general and regional anesthesia is influenced by several factors, including the extent of the surgery, patient preference, and the presence of any contraindications.

During the surgery, the anesthesiologist continuously monitors the patient's vital signs, including heart rate, blood pressure, oxygen saturation, and respiratory rate. Advanced monitoring techniques, such as electrocardiography (ECG), capnography (monitoring of carbon dioxide levels), and pulse oximetry, are employed to ensure the patient remains stable throughout the procedure.

The anesthesiologist also manages fluid and blood loss, administering intravenous fluids and blood products as necessary to maintain hemodynamic stability. This vigilant monitoring and management are critical in preventing complications and ensuring the patient's safety. Pain management is a central aspect of anesthesiology in abdominoplasty [41].

Effective pain control not only enhances patient comfort but also facilitates smoother recovery

and reduces the risk of postoperative complications such as deep vein thrombosis or respiratory issues. The anesthesiologist employs a multimodal approach to pain management, combining various pharmacological agents and techniques to achieve optimal pain relief. This approach may include the use of opioids, non-opioid analgesics, local anesthetics, and nerve blocks. Preemptive analgesia, administering pain relief before the onset of pain, is often utilized to mitigate postoperative pain more effectively.

One of the critical components of postoperative care in abdominoplasty is the management of pain and nausea. The anesthesiologist plays a vital role in this phase, ensuring that the transition from anesthesia to recovery is as smooth and comfortable as possible. Patients are monitored closely as they wake up from anesthesia, with pain and nausea managed through medications and supportive care. Effective postoperative pain management helps in early mobilization, which is crucial for reducing the risk of complications and promoting faster recovery.

Anesthesiology in abdominoplasty also involves preparing for potential complications and emergencies. The anesthesiologist must be equipped to handle adverse reactions to anesthesia, such as allergic reactions, respiratory difficulties, or cardiovascular instability. In rare cases, complications like malignant hyperthermia, a severe reaction to certain anesthetic agents, can occur, requiring immediate and expert intervention. The anesthesiologist's expertise in identifying and managing these complications is essential for ensuring patient safety.

Furthermore, the anesthesiologist collaborates closely with the surgical team to optimize the surgical environment and outcomes. This collaboration includes coordinating the timing of anesthetic induction and emergence with the surgical schedule, ensuring adequate muscle

relaxation for the procedure, and managing any intraoperative challenges that may arise. Effective communication and teamwork between the anesthesiologist and surgeon are vital for a successful abdominoplasty.

Patient education and communication are also important aspects of anesthesiology in abdominoplasty. Before the surgery, the anesthesiologist discusses the anesthetic plan with the patient, addressing any concerns and explaining the risks and benefits of different anesthetic options. This dialogue helps to build trust and reduce anxiety, ensuring that the patient feels informed and confident about the procedure. Postoperatively, the anesthesiologist continues to provide information and support, guiding the patient through the recovery process and addressing any issues that may arise.

In recent years, advancements in anesthetic techniques and medications have further enhanced the safety and efficacy of anesthesia in abdominoplasty. The development of shorter-acting anesthetic agents, improved monitoring technologies, and refined pain management strategies have contributed to better patient outcomes and a more comfortable surgical experience. These advancements reflect the ongoing evolution of the field of anesthesiology and its commitment to improving patient care.

In conclusion, anesthesiology is a critical component of abdominoplasty surgery, encompassing comprehensive preoperative assessment, meticulous intraoperative management, and effective postoperative care. The anesthesiologist's expertise ensures patient safety, comfort, and optimal surgical outcomes, making it an indispensable part of the abdominoplasty process. Through careful planning, vigilant monitoring, and advanced pain management techniques, anesthesiology contributes significantly to the success of this transformative surgical procedure. As the field continues to evolve, ongoing research and

innovation in anesthetic practices will further enhance the safety and effectiveness of abdominoplasty, ultimately improving the quality of care for patients undergoing this procedure.

Abdominoplasty Surgery Techniques

Abdominoplasty, often known as a "tummy tuck," is a popular cosmetic surgical procedure aimed at enhancing the appearance of the abdominal region. This procedure involves the removal of excess skin and fat, as well as the tightening of weakened or separated abdominal muscles, resulting in a smoother and firmer abdominal profile. Abdominoplasty has gained considerable traction over the years due to its significant impact on a patient's physical appearance and self-esteem. However, the success of this procedure is largely dependent on the surgical techniques employed. Various techniques have been developed and refined to cater to the diverse needs of patients, offering customized solutions that address specific aesthetic and functional concerns. This comprehensive exploration of abdominoplasty surgery techniques provides an in-depth understanding of the options available, their applications, and the outcomes they can achieve. The evolution of abdominoplasty techniques can be traced back to the early 20th century, with significant advancements occurring in the latter half of the century. Initially, abdominoplasty was primarily focused on the removal of excess skin and fat for patients with significant weight loss or postpartum changes. As surgical expertise and technology advanced, the procedure evolved to include muscle tightening and more sophisticated methods of skin excision and fat removal. Today, abdominoplasty encompasses a range of techniques designed to address specific issues, such as the standard full abdominoplasty, mini abdominoplasty, extended abdominoplasty, circumferential abdominoplasty, and the combination of

abdominoplasty with liposuction. Each technique offers unique benefits and is selected based on the patient's individual anatomy, goals, and the extent of correction required.

The most common form of abdominoplasty is the full abdominoplasty, which is suitable for patients with substantial excess skin and muscle laxity. This technique involves a horizontal incision made just above the pubic area, extending between the hip bones. Through this incision, the surgeon can access the underlying muscles and fascia to repair any separation or weakness, typically caused by pregnancy or significant weight fluctuations. The skin is then pulled down, excess skin is removed, and a new opening for the navel is created. This comprehensive approach not only tightens the abdominal wall but also significantly improves the overall contour and appearance of the abdomen. The full abdominoplasty is particularly beneficial for individuals who require extensive correction and desire a dramatic transformation.

For patients with minimal excess skin and localized fat deposits primarily below the navel, the mini abdominoplasty offers a less invasive alternative. This technique involves a shorter incision, typically confined to the lower abdomen, and does not usually require the repositioning of the navel. The surgeon tightens the lower abdominal muscles and removes a smaller amount of excess skin compared to a full abdominoplasty. Mini abdominoplasty is ideal for patients who are close to their ideal body weight but struggle with a small pouch or loose skin in the lower abdomen that is resistant to diet and exercise. This procedure offers quicker recovery times and fewer complications, making it an attractive option for those seeking more modest improvements.

Extended abdominoplasty expands on the traditional full abdominoplasty by addressing not only the abdominal region but also the flanks and lower back. This technique involves a longer

incision that extends around the hips, allowing for the removal of excess skin and fat from the sides and back in addition to the abdomen. This approach is particularly beneficial for patients who have undergone significant weight loss and have excess skin and fat deposits extending beyond the abdominal area. The extended abdominoplasty provides a more comprehensive body contouring solution, enhancing the silhouette from multiple angles and improving the overall balance and proportion of the body.

For patients who require even more extensive correction, the circumferential abdominoplasty, also known as a lower body lift, may be the most appropriate option. This technique involves an incision that encircles the entire waistline, allowing for the removal of excess skin and fat from the abdomen, flanks, and lower back, as well as the buttocks and thighs. This procedure is often indicated for patients who have experienced massive weight loss, resulting in significant skin laxity and drooping across multiple areas of the body. The circumferential abdominoplasty provides the most dramatic and transformative results, effectively addressing skin excess and improving the overall contour and proportion of the lower body.

In recent years, the combination of abdominoplasty with liposuction has gained popularity as an effective method for enhancing the results of the procedure. Liposuction involves the removal of localized fat deposits through small incisions using a suction device. When combined with abdominoplasty, liposuction can be used to sculpt and refine the abdominal contours, providing a smoother and more defined appearance. This combination approach is particularly beneficial for patients who have both excess skin and stubborn fat deposits that are resistant to diet and exercise. By integrating liposuction, surgeons can achieve more harmonious and natural-looking results,

enhancing the overall aesthetic outcome of the abdominoplasty.

Another advanced technique in abdominoplasty is the use of progressive tension sutures, which helps to secure the skin flap to the underlying tissues without the need for surgical drains. Traditionally, drains have been used to remove excess fluid and prevent complications such as seroma (fluid accumulation). However, the use of progressive tension sutures minimizes dead space, reducing the likelihood of fluid buildup and eliminating the need for drains. This innovation has been shown to improve patient comfort, reduce postoperative complications, and enhance the overall recovery experience.

The incorporation of advanced technologies such as laser-assisted liposuction and radiofrequency energy devices has further refined abdominoplasty techniques. Laser-assisted liposuction uses laser energy to liquefy fat before removal, resulting in less trauma to the surrounding tissues and promoting skin tightening. Radiofrequency energy devices deliver controlled thermal energy to stimulate collagen production and improve skin elasticity. These technologies can be used in conjunction with traditional abdominoplasty techniques to enhance skin tightening and improve the overall quality of the results.

The choice of abdominoplasty technique is influenced by various factors, including the patient's anatomy, the extent of correction required, and the surgeon's expertise. A thorough preoperative assessment and consultation are essential to determine the most appropriate approach for each individual patient. During this consultation, the surgeon evaluates the patient's medical history, physical condition, and aesthetic goals to develop a personalized surgical plan. This individualized approach ensures that the chosen technique aligns with the patient's expectations and provides the best possible outcome.

Postoperative care and recovery play a crucial role in the success of abdominoplasty surgery. Regardless of the technique employed, patients are typically advised to avoid strenuous activities and heavy lifting for several weeks following the procedure. Compression garments are often recommended to support the healing tissues, reduce swelling, and enhance the final contours. Pain management, wound care, and follow-up visits are integral components of the recovery process, ensuring that any complications are promptly addressed and that the healing progresses smoothly.

The psychological impact of abdominoplasty should not be underestimated. Many patients seek this procedure to address long-standing concerns about their abdominal appearance, which can significantly affect their self-esteem and quality of life. By restoring a more youthful and toned abdominal contour, abdominoplasty can boost confidence and improve overall well-being. However, it is important for patients to have realistic expectations and understand that while the procedure can provide significant improvements, it is not a substitute for a healthy lifestyle.

In conclusion, abdominoplasty surgery techniques have evolved significantly over the years, offering a range of options to address the diverse needs and goals of patients. From the traditional full abdominoplasty to the more specialized mini, extended, and circumferential techniques, each approach provides unique benefits tailored to individual patient requirements. The integration of advanced technologies and innovative techniques has further enhanced the safety, efficacy, and aesthetic outcomes of abdominoplasty. Through careful patient assessment, personalized surgical planning, and comprehensive postoperative care, surgeons can achieve transformative results that improve both physical appearance and psychological well-being. As the field of cosmetic surgery continues

to advance, ongoing research and innovation will undoubtedly lead to further refinements in abdominoplasty techniques, ensuring that patients receive the highest standard of care and the most satisfying results.

Conclusion

Abdominoplasty surgery techniques have significantly evolved, offering a variety of options to address the unique needs and goals of patients. From the comprehensive full abdominoplasty to the more focused mini abdominoplasty, and the extensive extended and circumferential procedures, each technique provides tailored solutions for achieving a firmer and more contoured abdominal profile. The integration of liposuction with abdominoplasty has further refined results, enabling more precise body sculpting and enhanced overall outcomes. Innovations such as progressive tension sutures and laser-assisted liposuction have improved the safety and efficacy of these procedures, reducing complications and promoting better skin tightening. A personalized approach, beginning with a thorough preoperative assessment, ensures that the chosen technique aligns with the patient's anatomy and aesthetic aspirations. Effective postoperative care, including pain management and the use of compression garments, is crucial for a smooth recovery and optimal results. As the field of cosmetic surgery continues to advance, ongoing research and the development of new technologies will likely lead to even more refined techniques, further enhancing patient satisfaction and the overall success of abdominoplasty. Ultimately, these advancements contribute to significant improvements in physical appearance, self-esteem, and quality of life for patients undergoing abdominoplasty.

References

- [1]SR Fazeli Daryasari, et al., The role of adiponectin in gestational diabetes mellitus, preeclampsia and obesity during pregnancy: A systematic review. *Iranian Journal of Endocrinology and Metabolism*. **2018**; 19(5): 370-83. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [2]F Pahlavan, et al., Role of Adiponectin Hormone in Fertility and Sterility: A Systematic Review. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. **2018**; 21(5):97-107. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [3]N Mobaraki Asl, et al., Comparing the effectiveness of intravenous oxytocin versus rectal misoprostol in the management of third-stage of labour after second-trimester abortion. *International Journal of Women's Health and Reproduction Sciences*. **2016**; 4(4):181-4. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [4]S Cuschieri, The STROBE guidelines. *Saudi journal of anaesthesia*. **2019**; 13(Suppl 1): S31. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [5]D Panidis, et al., Serum adiponectin levels in women with polycystic ovary syndrome. *Human reproduction*. **2003**; 18(9):1790-6. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [6]P-H Ducluzeau, et al. Glucose-to-insulin ratio rather than sex hormone-binding globulin and adiponectin levels is the best predictor of insulin resistance in nonobese women with polycystic ovary syndrome. *The Journal of Clinical Endocrinology & Metabolism*. **2003**; 88(8):3626-31. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [7]F Orio Jr, et al., Adiponectin levels in women with polycystic ovary syndrome. *The Journal of Clinical Endocrinology & Metabolism*. **2003** 88(6): 2619-23. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [8]J Spranger, et al. Adiponectin is independently associated with insulin sensitivity in women with polycystic ovary syndrome. *Clinical endocrinology*. **2004** 61(6):738-46. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [9]A Khosrowbeygi, H Ahmadvand, Serum levels of homocysteine, adiponectin and uric acid in gestational diabetes mellitus compared with normal pregnancy. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. **2017** 19(38):13-23. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [10] E Carmina, et al., Evidence for altered adipocyte function in polycystic ovary syndrome. *European Journal of Endocrinology*. **2005** 152(3):389-94. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [11] MSM Ardawi, AA Rouzi, Plasma adiponectin and insulin resistance in women with polycystic ovary syndrome. *Fertility and Sterility*. **2005** 83(6):1708-16. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [12] H Escobar-Morreale, et al. Adiponectin and resistin in PCOS: a clinical, biochemical and molecular genetic study. *Human reproduction*. **2006** 21(9):2257-65. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [13] NE Gulcelik, et al., Adiponectin is an independent determinant of insulin resistance in women with polycystic ovary syndrome. *Gynecological endocrinology*. **2006** 22(9):511-5. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [14] W Bik, et al., The relationship between metabolic status and levels of adiponectin and ghrelin in lean women with polycystic ovary syndrome. *Gynecological Endocrinology*. **2007** 23(6):325-31. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [15] V Aroda, et al., Circulating and cellular adiponectin in polycystic ovary syndrome: relationship to glucose tolerance and insulin action. *Fertility and sterility*. **2008** 89(5):1200-8. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [16] KA Toulis, et al. Adiponectin levels in women with polycystic ovary syndrome: a systematic review and a meta-analysis. *Human reproduction update*. **2009** 15(3):297-307. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

- [17] I Katsikis., et al., Oligo-ovulation or anovulation and hyperandrogenemia contribute to the decreased serum adiponectin levels in normal-weight women with PCOS with obesity and insulin resistance. *Fertility and Sterility*. **2009** 91(4): e3. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [18] M Yilmaz., et al. Serum resistin and adiponectin levels in women with polycystic ovary syndrome. *Gynecological Endocrinology*. **2009** 25(4):246-52. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [19] M Olszanecka-Glinianowicz., et al., Serum adiponectin and resistin in relation to insulin resistance and markers of hyperandrogenism in lean and obese women with polycystic ovary syndrome. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. **2011** 154(1):51-6. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [20] FV Comim., et al., Adiponectin and its receptors in the ovary: further evidence for a link between obesity and hyperandrogenism in polycystic ovary syndrome. *PloS one*. **2013** 8(11): e80416. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [21] T Artimani., et al. Downregulation of adiponectin system in granulosa cells and low levels of HMW adiponectin in PCOS. *Journal of assisted reproduction and genetics*. **2016** 33(1):101-10. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [22] MA Alfaqih., et al., Lower levels of serum adiponectin and the T allele of rs1501299 of the ADIPOQ gene are protective against polycystic ovarian syndrome in Jordan. *Korean journal of family medicine*. **2018** 39(2):108. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [23] DP Baldani., et al., Altered leptin, adiponectin, resistin and ghrelin secretion may represent an intrinsic polycystic ovary syndrome abnormality. *Gynecological Endocrinology*. **2019** 35(5):401-5. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [24] DM Bannigida., SB Nayak SB. Serum visfatin and adiponectin-markers in women with polycystic ovarian syndrome. *Archives of physiology and biochemistry*. **2020** 126(4):283-6. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [25] X Sun., et al., Family-based association study of rs17300539 and rs12495941 polymorphism in adiponectin gene and polycystic ovary syndrome in a Chinese population. *Medical science monitor: international medical journal of experimental and clinical research*. **2017** 23:78. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [26] NS Cardoso., et al. Polycystic ovary syndrome associated with increased adiposity interferes with serum levels of TNF-alpha and IL-6 differently from leptin and adiponectin. *Archives of Endocrinology and Metabolism*. **2020** 64(1):4-10. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [27] S Shorakae., et al. High-molecular-weight adiponectin is inversely associated with sympathetic activity in polycystic ovary syndrome. *Fertility and Sterility*. **2018** 109(3):532-9. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [28] Jiang S, Zhang Y, Alsaikhan F, Jalil AT, Gol MK, Tarighatnia A. A meta-analysis review of the effect of Zn-doped synthetic polymer materials on bone regeneration. *Journal of Drug Delivery Science and Technology*. **2022** 1; 76:103792. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [29] Gol MK, Payami S, Lotfi A. Study of the effect of ear acupressure on stress and serum cortisol level before rhinoplasty surgery: a randomized clinical trial. *Crescent Journal of Medical & Biological Sciences*. **2020** 1;7(2). [[Google Scholar](#)], [[Publisher](#)]
- [30] Khanbabayi Gol M, Arefi N, Jafari M, Farzin H, Aghamohammadi D. Prevalence of Port-Related Infections and Their Predisposing Factors in Women with Breast Cancer under Chemotherapy. *Iranian Quarterly Journal of*

Breast Disease. **2018** 10;11(2):7-15. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[31] Piroozkhah, Moein, et al. "Guanylate cyclase-C Signaling Axis as a theragnostic target in colorectal cancer: a systematic review of literature." *Frontiers in Oncology*, **2023** 13. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[32] AZHOUGH, RAMIN, et al. "Surgical Outcome and Complications of Lateral Internal Sphincterotomy in Chronic Anal Fissure-A Cross-sectional Study." *Journal of Clinical & Diagnostic Research*, **2021** 15.12. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[33] Jalali, Pooya, et al. "UBE2C: A pan-cancer diagnostic and prognostic biomarker revealed through bioinformatics analysis." *Cancer Reports*, **2024** 7(4): e2032. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[34] Azhough, Ramin, et al. "Pilonidal sinus surgery with mechanical drilling technique." *Indian Journal of Surgery*, **2020** 82: 824-827. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[35] Azhough, Ramin, et al. "Intradermal methylene blue analgesic application in posthemorrhoidectomy pain management: a randomized controlled trial." *Frontiers in Surgery*, **2024** 11: 1354328. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[36] K Hadavand Mirzaei, et al., Application of Robotic Instruments in Hip Arthroplasty

Surgery Based on Practical Tips a Systematic Review, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2024** 3 (2), 593-609 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[37] S Sourili, What are the radiological criteria for identifying the Corona virus in CT scan images?, *Eurasian Journal of Chemical, Medicinal and Petroleum Research (EJCMPR)*, **2024** 3(2), 577-593 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[38] S Soubkari, Covid CT Chest Finding, *Eurasian Journal of Chemical, Medicinal and Petroleum Research (EJCMPR)*, **2024** 3(2), 563-576 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[39] S Asadimehr, Examining Drilling Problems and Practical Solutions Regarding them, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2024** 3 (2), 552-562 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[40] S Asadimehr, Investigating the Use of Drilling Mud and the Reasons for its Use, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2024** 3 (2), 543-551 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[41] S Soubhak, Nursing actions in patients undergoing laryngectomy in relation to reducing anxiety and depression, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, **2024** 3 (2), 530-542 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]