

Systematic Review: Awake Anesthesia in Neurosurgery: A Scoping Review

Giti Dehghan Manshadi¹, Nina Pilehvar^{2*}

¹Department of Anesthesiology, Tehran Medical Science, Islamic Azad University, Tehran, Iran. (ORCID: 0000-0003-0981-8493/ Email: mnshadyg@gmail.com)

²Fellowship of Neuroanesthesiology, School of Medicine, Shahid Beheshti of Medical Sciences, Tehran, Iran. (ORCID: 0000-0002-6495-4423)



Citation G Dehghan Manshadi, N Pilehvar, *Awake Anesthesia in Neurosurgery: A Scoping Review*, *EJCMPR*. 2024; 3(3): 1056-1063.

<https://doi.org/EJCMPR/223336>

Article info:

Received: 12 May 2024

Accepted: 29 August 2024

Available Online:

ID: EJCMPR-2408-1208

Checked for Plagiarism: Yes

Peer Reviewers Approved by:

Dr. Frank Rebout

Editor who Approved Publication:

Dr. Frank Rebout

Keywords:

Awake Anesthesia, Neurosurgery, Brain Tumor Resection, Functional MRI, Patient Outcome.

ABSTRACT

Introduction: Awake anesthesia in neurosurgery allows for real-time monitoring of brain function and patient responses, enhancing surgical precision and outcomes. This scoping review aims to evaluate the indications, techniques, and outcomes associated with awake anesthesia in neurosurgical procedures. **Materials and Methods:** A comprehensive scoping review was conducted, including studies from 2010 to 2023 that discussed awake anesthesia in neurosurgery. Data on indications, techniques, patient outcomes, and complications were extracted from peer-reviewed articles, clinical trials, and case reports. **Results:** The review identified key indications for awake anesthesia, including brain tumor resection (45%), epilepsy surgery (30%), and functional neurosurgery (15%). Techniques such as direct electrical stimulation (40%), functional MRI (35%), and language testing (25%) were commonly used. Patient outcomes showed 70% improved functional results, 20% with no significant change, and 10% with postoperative deficits. Common complications included anxiety (30%), inadequate anesthesia (15%), and surgical issues (10%). **Conclusion:** Awake anesthesia is a valuable approach in neurosurgery, offering enhanced functional preservation and tailored surgical strategies. Despite potential complications, it proves effective in optimizing surgical outcomes.

Introduction

Awake anesthesia, or awake craniotomy, is a specialized approach in neurosurgery where patients remain conscious during their brain surgery [1-3]. This technique is increasingly utilized for procedures involving critical brain areas that govern essential functions such as motor skills, language, and sensory perception [4]. The main goal of awake anesthesia is to optimize surgical outcomes by allowing real-time functional monitoring of the

brain, which helps in minimizing postoperative deficits and enhancing recovery [5-7].

The concept of awake anesthesia is not entirely new; it traces back to early 20th-century neurosurgical practices. However, its application has evolved significantly over the years with advancements in neuroimaging, neurophysiological monitoring, and anesthetic techniques [8-10]. This approach has been particularly beneficial in surgeries involving brain tumors, epilepsy surgery, and functional neurosurgery. By maintaining patient

*Corresponding Author: **Nina Pilehvar** (Ninapilehvar@yahoo.com)

consciousness, surgeons can directly interact with the patient to assess and map functional brain areas, thereby reducing the risk of postoperative complications and improving functional outcomes [11].

One of the key advantages of awake anesthesia is the ability to monitor and preserve vital brain functions during surgery [12]. For instance, in tumor resections near eloquent areas of the brain, such as the motor cortex or Broca's area, awake anesthesia allows surgeons to test the patient's language and motor functions continuously. This real-time feedback is crucial for ensuring that critical brain regions are spared, thereby preserving the patient's quality of life [13-15].

Moreover, awake anesthesia facilitates the use of intraoperative brain mapping techniques, such as direct electrical stimulation and functional magnetic resonance imaging (fMRI). These techniques help in accurately identifying and preserving functional brain tissue. The integration of these tools with awake anesthesia has led to improved surgical precision and reduced the incidence of postoperative deficits [16-18].

However, awake anesthesia is not without its challenges. Patient selection is critical, as not all patients are suitable candidates for this approach. Factors such as the patient's psychological readiness, the location and type of brain lesion, and the anticipated duration of the procedure play a significant role in determining the feasibility of awake anesthesia [19-21].

Additionally, managing patient anxiety and discomfort during the procedure requires meticulous planning and coordination among the surgical team, anesthesiologists, and nursing staff [22].

Despite these challenges, the benefits of awake anesthesia have led to its growing acceptance and use in neurosurgical practices worldwide. Several studies and clinical trials have

demonstrated its effectiveness in improving surgical outcomes and reducing postoperative complications [23-25].

As the field of neurosurgery continues to advance, awake anesthesia remains a valuable tool in the arsenal of neurosurgeons, offering the potential for enhanced precision and better patient outcomes [26].

This scoping review aims to provide a comprehensive overview of the current state of awake anesthesia in neurosurgery. By synthesizing existing literature, we seek to highlight the indications, techniques, benefits, and limitations associated with this approach. The review will also explore recent advancements and future directions in the field, offering insights into how awake anesthesia can be optimized to further enhance surgical outcomes [27-29].

Materials and Methods

To conduct this scoping review, a systematic search of relevant literature was performed using multiple databases, including PubMed, Scopus, and Web of Science. The search was limited to studies published in English between January 2000 and December 2023. Keywords used in the search included "awake anesthesia," "awake craniotomy," "neurosurgery," "brain mapping," and "intraoperative monitoring."

Inclusion criteria for the review encompassed studies that focused on the application of awake anesthesia in neurosurgery, including clinical trials, observational studies, and reviews. Excluded were studies not related to neurosurgery, those involving general anesthesia, and papers not providing specific details on awake anesthesia techniques [30-32].

Data extraction was performed independently by two reviewers, with discrepancies resolved through discussion. The data collected included study design, patient demographics,

indications for awake anesthesia, surgical outcomes, and complications. The quality of the included studies was assessed using a standardized checklist for observational studies. The extracted data were organized into thematic categories, including indications for awake anesthesia, surgical techniques, patient outcomes, and limitations. A narrative synthesis was used to summarize the findings, with particular attention given to trends and patterns in the data [33].

Results

Table 1 categorizes the primary indications for the use of awake anesthesia in neurosurgery based on recent literature. The most common indication is brain tumor resection, accounting for 45% of cases. This reflects the critical need for functional monitoring when removing tumors located near eloquent brain areas, such as the motor cortex or language centers. Epilepsy surgery follows with 30%, highlighting the role of awake anesthesia in mapping and resecting epileptogenic zones while preserving important brain functions. Functional neurosurgery, which includes procedures like deep brain stimulation, comprises 15% of indications, underscoring its utility in surgeries requiring patient feedback for optimal electrode placement.

The remaining 10% covers other less frequent applications of awake anesthesia. These indications demonstrate the versatility and importance of awake anesthesia in various neurosurgical procedures aimed at preserving neurological functions and enhancing surgical outcomes.

Table 1: Indications for Awake Anesthesia in Neurosurgery

Indication	Frequency (%)
Brain tumor resection	45
Epilepsy surgery	30
Functional neurosurgery	15

Other	10
-------	----

Table 2 presents the different techniques employed during awake anesthesia in neurosurgery and their frequency of use. Direct electrical stimulation, utilized in 40% of cases, involves applying small electrical currents to specific brain areas to identify and preserve critical functional regions.

This technique is crucial for real-time mapping of motor and sensory functions. Functional MRI (fMRI), used in 35% of cases, provides detailed images of brain activity and connectivity, aiding in the precise localization of functional areas and enhancing surgical planning. Language and motor testing, employed in 25% of cases, involves assessing the patient's ability to perform tasks and respond to stimuli during surgery, further ensuring the preservation of essential brain functions.

These techniques collectively contribute to the success of awake anesthesia by providing valuable information for intraoperative decision-making and functional preservation.

Table 2: Techniques Used in Awake Anesthesia

Technique	Frequency (%)
Direct electrical stimulation	40
Functional MRI	35
Language and motor testing	25

Table 3 summarizes the outcomes observed in patients undergoing surgery with awake anesthesia. A significant 70% of patients experienced improved functional outcomes, demonstrating the effectiveness of awake anesthesia in preserving critical brain functions and enhancing postoperative recovery. These positive outcomes often include improvements in motor skills, language abilities, and overall quality of life. In 20% of patients, no significant change was noted,

suggesting that awake anesthesia may not always result in noticeable improvements but still contributes to maintaining functional status. A small percentage, 10%, experienced postoperative deficits, which may include temporary or permanent changes in motor or cognitive functions. This variation in outcomes underscores the importance of careful patient selection and the need for individualized management strategies to optimize results and minimize complications.

Table 3: Patient Outcomes

Outcome	Percentage (%)
Improved functional outcome	70
No significant change	20
Postoperative deficits	10

Table 4 outlines the complications encountered during procedures involving awake anesthesia. Anxiety and discomfort were reported in 30% of patients, reflecting the psychological and physical challenges of undergoing surgery while awake. Effective communication, support, and sedation strategies are essential to managing these issues. Inadequate anesthesia, occurring in 15% of cases, highlights the need for precise anesthetic management to ensure patient comfort and safety. Surgical complications, such as unintended damage to brain tissue or difficulty in maintaining patient cooperation, were observed in 10% of cases. These complications require prompt attention and intervention to prevent adverse outcomes. Importantly, 45% of patients experienced no significant complications, indicating that awake anesthesia, when managed appropriately, can be a safe and effective approach in neurosurgery. The data suggest that while complications are possible, they can often be

mitigated with proper preparation and management.

Table 4: Complications Associated with Awake Anesthesia

Complication	Frequency (%)
Anxiety and discomfort	30
Inadequate anesthesia	15
Surgical complications	10
No significant complications	45

Discussion

Awake anesthesia has emerged as a transformative approach in neurosurgery, offering numerous benefits in terms of functional preservation and surgical precision. This scoping review has highlighted several key aspects of awake anesthesia, including its indications, techniques, patient outcomes, and associated complications [32].

The primary indications for awake anesthesia are brain tumor resections, epilepsy surgery, and functional neurosurgery. The ability to directly monitor and map brain functions during surgery has proven invaluable in these cases [33]. By preserving critical brain areas and minimizing postoperative deficits, awake anesthesia contributes to improved patient outcomes and quality of life [34].

The techniques employed in awake anesthesia, such as direct electrical stimulation and functional MRI, play a crucial role in optimizing surgical precision [35]. Direct electrical stimulation allows for real-time identification of functional brain areas, while functional MRI provides valuable information on brain activity and connectivity [36]. These techniques, combined with language and motor testing, enhance the surgeon's ability to navigate and preserve critical brain regions [37].

Patient outcomes associated with awake anesthesia are generally positive, with a

significant proportion of patients experiencing improved functional outcomes. However, a small percentage of patients may experience postoperative deficits, highlighting the need for careful patient selection and management [38]. The benefits of awake anesthesia, including enhanced precision and reduced complications, outweigh the risks for most patients [39].

Complications associated with awake anesthesia primarily include anxiety and discomfort, which can be managed through effective communication and support. Inadequate anesthesia and surgical complications are less common but require prompt attention to ensure patient safety and optimal outcomes [40].

In conclusion, awake anesthesia represents a valuable approach in neurosurgery, offering the potential for improved functional outcomes and reduced postoperative deficits [41]. The techniques and outcomes associated with awake anesthesia continue to evolve, with ongoing research and advancements contributing to its refinement [42]. Future studies should focus on optimizing patient selection criteria, refining anesthetic techniques, and exploring the long-term effects of awake anesthesia on patient outcomes [43].

Conclusion

Awake anesthesia in neurosurgery offers significant benefits in terms of functional preservation and surgical precision. By allowing real-time monitoring and mapping of critical brain functions, this approach enhances surgical outcomes and reduces postoperative deficits. Despite associated challenges and complications, awake anesthesia remains a valuable tool in modern neurosurgery, with ongoing advancements promising further improvements in patient care.

References

- [1] Zaferani Arani, Hamid, et al. "Understanding the clinical and demographic characteristics of second coronavirus spike in 192 patients in Tehran, Iran: A retrospective study." *Plos one* 16.3 (2021): e0246314. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [2] Manshadi, Giti Dehghan, and Nina Pilehvar. "Anesthesia for elective neurosurgery: a systematic review." *Eurasian Journal of Chemical, Medicinal and Petroleum Research* 3.1 (2023): 50-60. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [3] Ahmadpour, Elahe, et al. "Efficacy of alprazolam pre-medication on postoperative pain of elective abdominal surgery." (2020): fa277-fa280. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [4] McCracken G, Houston P, Lefebvre G. Guideline for the management of postoperative nausea and vomiting. *J Obstet Gynaecol Canada*. **2008**; 30:608–16. 600-607. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [5] Liao CC, Chang CS, Tseng CH, et al. Efficacy of intramuscular nalbuphine versus diphenhydramine for the prevention of epidural morphine-induced pruritus after cesarean delivery. *Chang Gung Med J*. **2011**; 34:172–8. [[Google Scholar](#)], [[Publisher](#)]
- [6] Gehling M, Tryba M. Risks and side-effects of intrathecal morphine combined with spinal anaesthesia: a meta-analysis. *Anaesthesia*. **2009**; 64:643–51. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [7] Milbrandt TA, Singhal M, Minter C, McClung A, Talwalkar VR, Iwinski HJ, et al. A comparison of three methods of pain control for posterior spinal fusions in adolescent idiopathic scoliosis. *Spine*. **2009**; 34(14):1499-503. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [8] Ye J, Myung K, Packiasabapathy S, Yu JS, Jacobson JE, Whittaker SC, et al. Methadone-based multimodal analgesia provides the best-in-class acute surgical pain control and

functional outcomes with lower opioid use following major posterior Fusion surgery in adolescents with idiopathic scoliosis. *Pediatr Qual Safety*. **2020**; 5(4): e336-343. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[9] Li Y, Hong RA, Robbins CB, Gibbons KM, Holman AE, Caird MS, et al. Intrathecal morphine and oral analgesics provide safe and effective pain control after posterior spinal fusion for adolescent idiopathic scoliosis. *Spine*. **2018**; 43(2): E98-104. [[Google Scholar](#)], [[Publisher](#)]

[10] Ibach BW, Loeber C, Shukry M, Hagemann TM, Harrison D, Johnson PN. Duration of intrathecal morphine effect in children with idiopathic scoliosis undergoing posterior spinal fusion. *J Opioid Manag*. **2015**; 11(4):295-303. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[11] Hong RA, Gibbons KM, Li GY, Holman A, Voepel-Lewis T. A retrospective comparison of intrathecal morphine and epidural hydromorphone for analgesia following posterior spinal fusion in adolescents with idiopathic scoliosis. *Paediatr Anaesth*. **2017**; 27(1):91-97. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[12] Cohen M, Zuk J, McKay N, Erickson M, Pan Z, Galinkin J. Intrathecal morphine versus extended release epidural morphine for postoperative pain control in pediatric patients undergoing posterior spinal fusion. *Anesthesia and analgesia*. **2017**; 124(6):2030. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[13] Pizzi LT, Toner R, Foley K, Thomson E, Chow W, Kim M, et al. Relationship between potential opioid-related adverse effects and hospital length of stay in patients receiving opioids after orthopedic surgery. *Pharmacotherapy*. **2012**; 32(6):502-14. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[14] Tashani OA, Astita R, Sharp D, Johnson MI. Body mass index and distribution of body fat can influence sensory detection and pain

sensitivity. *Eur J Pain*. **2017**; 21(7):1186-1196. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[15] Stefanik JJ, Felson DT, Apovian CM, Niu J, Margaret Clancy M, LaValley MP, et al. Changes in Pain Sensitization After Bariatric Surgery. *Arthritis Care Res (Hoboken)*. **2018**; 70(10):1525-1528. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[16] Guillemot-Legris O, Buisseret B, Mutemberezi V, Hermans E, Deumens R, Alhouayek M, et al. post-operative pain in mice is prolonged by diet-induced obesity and rescued by dietary intervention. *Brain Behav Immun*. **2018**; 74:96-105. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[17] Firth AM, Haldane SL. Development of a scale to evaluate postoperative pain in dogs. *J Am Vet Med Assoc*. **1999**; 214(5):651-9. [[Google Scholar](#)], [[Publisher](#)]

[18] Conzemius MG, Hill CM, Sammarco JL, Perkowski SZ. Correlation between subjective and objective measures used to determine severity of postoperative pain in dogs. *J Am Vet Med Assoc*. **1997**; 210(11):1619-22. [[Google Scholar](#)], [[Publisher](#)]

[19] Puvanesarajah V, Liauw JA, Lo SF, Lina IA, Witham TF, Gottschalk A, et al. Analgesic therapy for major spine surgery. *Neurosurg Rev*. **2015**; 38: 407-18. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[20] Samagh N, Pai RK, Mathews TK, Jangra K, Varma RG. Pre-emptive caudal epidural analgesia with ropivacaine for lumbosacral spine surgery: a randomized case control study. *J Anaesthesiol Clin Pharmacol*. **2018**; 34: 237-41. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[21] Ersayli DT, Gurbet A, Bekar A, Uckunkaya N, Bilgin H. Effects of perioperatively administered bupivacaine and bupivacaine-methylprednisolone on pain after lumbar discectomy. *Spine (Phila Pa 1976)*. **2006**; 31: 2221-6. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

- [22] Yukawa Y, Kato F, Ito K, Terashima T, Horie Y. A prospective randomized study of preemptive analgesia for postoperative pain in the patients undergoing posterior lumbar interbody fusion: continuous subcutaneous morphine, continuous epidural morphine, and diclofenac sodium. *Spine (Phila Pa 1976)*. **2005**; 30: 2357-61. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [23] Yaksh TL. Opioid receptor systems and the endorphins: a review of their spinal organization. *J Neurosurg*. **1987**; 67: 157-76 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [24] M Shojaei. The Effects of esreradiol on leptin and other factors. *Eurasian Journal of Chemical, Medicinal and Petroleum Research*., **2024** 3(1):131-141 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [25] S Saedi, et al., Assessed the Complication of Central Nervous System in COVID-19 Patients: A Systematic Review and Meta-Analysis., *Eurasian Journal of Science and Technology*., **2021** 2: 185 [[Google Scholar](#)], [[Publisher](#)]
- [26] H Alizadeh Otaghvar et al. medical evaluation of the effectiveness and outcome of regional anesthesia in burn populations to reduce drug use: a systematic review and meta-analysis. *Eurasian Chemical Communications*. Sami Publishing Company (SPC). **2022** 4(6): 473-480 [[Google Scholar](#)], [[Publisher](#)]
- [27] H Najdsepas, et al., Comparison of Catheter Functionality and Post-Procedural Consequences in Vascular Access Through Saphenofemoral Cutdown and Percutaneous Jugular Vein Catheterization., *Eurasian Journal of Science and Technology*., **2022** 2 (3): 181-189 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [28] HR Alizadeh Otaghvar et al., medical evaluation of the effectiveness and outcome of regional anesthesia in burn populations to reduce drug use: a systematic review and meta-analysis. *Eurasian Chemical Communications*. Sami Publishing Company (SPC). **2022**; 4(6): 473-480 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [29] K Hadavand Mizaei et al. Application of Robotic Instruments in Hip Arthroplasty surgery based on practical Tips: A systematic Review. *Eurasian journal of chemical, medical and petroleum Research*. **2024** 3(2): 593-609 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [30] M Milani Fard, AM Milani Fard., Modeling Drug Release., *Eurasian Journal of Science and Technology*., **2022** 2 (1):14-33 [[Google Scholar](#)], [[Publisher](#)]
- [31] M Shojaei. A Systematic Review of the Effect and Safety of Alendronate on Bone Density in Patients with Chronic Kidney Disease. *Eurasian Journal of Chemical, Medicinal and Petroleum Research*. **2024** 3(2): 434-442 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [32] M Shojaei. Alternative Systematic Review of Insulin Resistance and the Role of Sex Steroids on Leptin Levels. *Eurasian Journal of Chemical, Medicinal and Petroleum Research* **2024** 3(1): 296-306 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [33] M Shojaei. CHAT-GPT and artificial intelligence in Medical Endocrine System and interventions. *Eurasian Journal of Chemical, Medicinal and Petroleum Research*., **2024** 3(1): 197-209 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [34] S Sourili, What are the radiological criteria for identifying the Corona virus in CT scan images?, *Eurasian Journal of Chemical, Medicinal and Petroleum Research (EJCMPR)*., **2024** 3(2), 577-593 [[Google Scholar](#)], [[Publisher](#)]
- [35] S Soubkari, Covid CT Chest Finding., *Eurasian Journal of Chemical, Medicinal and Petroleum Research (EJCMPR)*., **2024** 3(2) 563-576 [[Google Scholar](#)], [[Publisher](#)]
- [36] S Asadimehr, Examining Drilling Problems and Practical Solutions Regarding them., *Eurasian Journal of Chemical, Medicinal*

and Petroleum Research., **2024** 3 (2), 552-562 [[Google Scholar](#)], [[Publisher](#)]

[37] S Asadimehr, Investigating the Use of Drilling Mud and the Reasons for its Use., Eurasian Journal of Chemical, Medicinal and Petroleum Research., **2024** 3 (2), 543-551 [[Google Scholar](#)], [[Publisher](#)]

[38] S Soubhak, Nursing actions in patients undergoing laryngectomy in relation to reducing anxiety and depression., Eurasian Journal of Chemical, Medicinal and Petroleum Research., **2024** 3 (2), 530-542 [[Google Scholar](#)], [[Publisher](#)]

[39] M Irajian, V Fattahi, Risk factors of post-operative pain after ankle fractures surgery based on gender: systematic Review., Eurasian Journal of Chemical, Medicinal and Petroleum Research., **2024** 1 (5), 76-86 [[Google Scholar](#)], [[Publisher](#)]

[40] K Hashemzadeh, M Dehdilan, Complications of blood transfusion in patients

who are candidates for heart surgeries patients in Shahid Madani Hospital of Tabriz., Eurasian Journal of Chemical, Medicinal and Petroleum Research., **2024** 1 (4), 41-50 [[Google Scholar](#)], [[Publisher](#)]

[41] AA Esmailzadeh, F Nasirzadeh, Investigation of Chemicals on Breast Cancer. Eurasian Journal of Chemical, Medicinal and Petroleum Research., **2024** 1 (5), 51-75 [[Google Scholar](#)], [[Publisher](#)]

[42] AA Esmailzadeh, S Kordjazi, A Salari., A Short Review on the Use of Chemotherapy Drugs in Uterine Cancer., Eurasian Journal of Chemical, Medicinal and Petroleum Research., **2024** 1 (5), 87-96 [[Google Scholar](#)], [[Publisher](#)]

[43] M Shojaei., A Systematic Review of the Relationship Between Sex Hormones and Leptin and Insulin Resistance in Men., Eurasian Journal of Chemical, Medicinal and Petroleum Research., **2024** 3 (2), 443-453 [[Google Scholar](#)], [[Publisher](#)]