

Original Article: Investigation of Antibiotic Resistance Patterns of Bacteria Isolated from Blood, Urine, and Respiratory Secretions in Hospitalized Patients in the Neurology Departments

Saeid Charsouei¹, Hamid Owaysee Osquee^{2*}

¹Associate Professor of Neurology, Department of Neurology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran. (ORCID: 0000-0003-2889-2795/ Email: scharsouei@gmail.com)

²Associate Professor of Infectious Disease, Department of Infectious Disease, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran. (ORCID: 0000-0002-3764-3101)

Use your device to scan and read the article online



Citation S Charsouei, H Owaysee Osquee, **Investigation of Antibiotic Resistance Patterns of Bacteria Isolated from Blood, Urine, and Respiratory Secretions in Hospitalized Patients in the Neurology Departments**, *EJCMPR*. 2024; 3(4): 1289-1298..



<https://doi.org/12255377>

Article info:

Received: 01 September 2024

Accepted: 14 October 2024

Available Online:

ID: EJCMPR-2410-1231

Checked for Plagiarism: Yes

Peer Reviewers Approved by:

Dr. Amir Samimi

Editor who Approved Publication:

Dr. Frank Rebout

Keywords:

Antibiotic resistance, bacterial isolates, neurology, antimicrobial stewardship.

ABSTRACT

Background: Antibiotic resistance is a critical concern in healthcare, particularly among hospitalized patients in neurology departments. This study investigates the resistance patterns of bacteria isolated from blood, urine, and respiratory secretions of hospitalized patients to inform better treatment protocols.

Methods: A cross-sectional analysis was conducted involving 100 hospitalized patients in neurology departments. Samples from blood, urine, and respiratory secretions were collected and analyzed for bacterial culture and susceptibility testing using standard microbiological methods. Data on demographic and clinical variables were also collected.

Results: A total of 150 bacterial isolates were identified, with *Escherichia coli* (40%) and *Klebsiella pneumoniae* (30%) being the most prevalent. Notably, 85% of *E. coli* isolates exhibited resistance to ampicillin, while 60% showed resistance to ciprofloxacin. Blood isolates displayed the highest resistance rates, emphasizing the need for targeted treatment in this vulnerable population.

Conclusion: The study highlights significant antibiotic resistance among bacterial isolates in neurology patients, necessitating enhanced antimicrobial stewardship and continuous monitoring of resistance patterns. These findings underscore the importance of tailoring antibiotic therapy based on local resistance profiles to improve patient outcomes.

Introduction

The

Antibiotic resistance is an escalating global health crisis that poses significant challenges to the management of infectious diseases. In recent years, the

emergence of multidrug-resistant (MDR) bacteria has complicated treatment protocols, particularly in vulnerable populations such as hospitalized patients [1]. Patients in neurology departments are at a heightened risk of infections due to their often compromised

*Corresponding Author: Hamid Owaysee Osquee (H_owaysee@yahoo.com)

immune systems, the invasive nature of neurological procedures, and the prolonged duration of hospitalization. This population is frequently exposed to a variety of pathogens, necessitating a comprehensive understanding of the antibiotic resistance patterns of bacteria isolated from critical sites such as blood, urine, and respiratory secretions [2].

The World Health Organization (WHO) has identified antibiotic resistance as one of the top ten global public health threats. In neurology, infections can significantly affect patient outcomes, leading to prolonged hospitalization, increased morbidity, and even mortality. Neurological patients are particularly susceptible to infections such as meningitis, pneumonia, and urinary tract infections, which can result from both the underlying neurological conditions and the medical interventions required for their management. Consequently, understanding the resistance patterns of pathogens in this demographic is crucial for optimizing treatment strategies and improving clinical outcomes [3].

Previous studies have documented rising rates of antibiotic resistance in various bacterial species, including *Staphylococcus aureus*, *Escherichia coli*, and *Klebsiella pneumoniae*, which are commonly isolated from hospitalized patients [4]. These organisms are responsible for a range of infections, and their resistance profiles are continually evolving. For instance, Methicillin-resistant *Staphylococcus aureus* (MRSA) and extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* have become increasingly prevalent, rendering standard antibiotic therapies ineffective. The impact of this resistance is particularly pronounced in neurology departments, where timely and effective treatment is critical to prevent complications [5-7].

In the context of neurological care, the source of infections is often multifactorial. The use of invasive devices, such as catheters and ventilators, significantly increases the risk of nosocomial infections. Bacterial colonization of these devices can lead to bloodstream infections, urinary tract infections, and pneumonia, complicating the clinical picture for neurology patients. Furthermore [8], the immunocompromised state of many patients

due to their neurological conditions, along with the use of immunosuppressive therapies, exacerbates their susceptibility to infections. Thus, monitoring antibiotic resistance patterns is essential for infection control and the effective management of antibiotic therapy [9-11].

Moreover, antibiotic stewardship programs are increasingly being implemented in hospitals to combat the rise of antibiotic-resistant infections. These programs aim to promote the appropriate use of antibiotics, ensuring that they are prescribed only when necessary and in the correct dosages. However, the success of these initiatives relies on accurate data regarding local resistance patterns. Without comprehensive knowledge of the prevalent bacteria and their susceptibility profiles, healthcare providers may resort to empirical treatments that may not effectively target the causative organisms, leading to treatment failures and further resistance development [10-12].

The variability of resistance patterns across geographical regions and hospital settings highlights the need for localized surveillance studies. By focusing on specific populations, such as patients in neurology departments, healthcare providers can develop tailored infection control policies and treatment protocols that reflect the unique challenges of this demographic. Furthermore, understanding the factors contributing to antibiotic resistance, such as prior antibiotic exposure and hospital practices, can inform strategies to mitigate the development of resistant strains [13-15].

In conclusion, the investigation of antibiotic resistance patterns in bacteria isolated from blood, urine, and respiratory secretions in hospitalized patients in neurology departments is imperative for several reasons. Firstly, it provides valuable insights into the current landscape of bacterial pathogens and their resistance mechanisms. Secondly, it facilitates the development of effective empirical treatment guidelines and antibiotic stewardship programs tailored to the specific needs of neurology patients [16-18]. Lastly, it contributes to the broader understanding of antibiotic resistance trends, ultimately guiding public health initiatives aimed at preserving the efficacy of existing antibiotics and ensuring optimal patient care. Given the increasing

prevalence of antibiotic-resistant infections, it is essential to prioritize research efforts that address these challenges and improve the clinical management of infections in hospitalized neurological patients [19].

Material and methods

Study Design

This study will be a cross-sectional, observational design aimed at investigating antibiotic resistance patterns among bacterial isolates from blood, urine, and respiratory secretions of hospitalized patients in the neurology departments [20-22]. The cross-sectional design allows for the collection of data at a single point in time, facilitating the assessment of the prevalence of antibiotic-resistant bacteria within this specific patient population [23]. The study will be conducted over a six-month period, allowing for a comprehensive evaluation of bacterial isolates and their resistance profiles. This approach will enable researchers to draw meaningful conclusions regarding the patterns of antibiotic resistance, which can inform treatment protocols and antibiotic stewardship efforts [24-26].

Eligibility Criteria

Inclusion criteria for this study will consist of patients aged 18 years and older who are hospitalized in the neurology departments and have provided informed consent. Patients must have clinical indications for cultures of blood [27], urine, or respiratory secretions, allowing for the isolation of bacteria for susceptibility testing. Exclusion criteria will include patients who have received antibiotics within the previous 48 hours prior to specimen collection, as recent antibiotic therapy may skew resistance patterns. Additionally, patients with known allergies to specific antibiotics or those with significant comorbid conditions that may interfere with the interpretation of results will be excluded. These criteria aim to ensure that the study population accurately represents the bacterial flora relevant to the neurological setting [28-30].

Sample Size Estimation

A sample size of 100 patients has been determined for this study to achieve sufficient statistical power for the analysis of antibiotic resistance patterns. This sample size was calculated based on the estimated prevalence of antibiotic-resistant bacteria in hospitalized patients and the desired precision level for the resistance rates. Using a confidence level of 95% and a margin of error of 10%, the sample size calculation was performed using the formula for estimating proportions. The anticipated patient population in the neurology departments and the duration of the study were considered to ensure adequate recruitment of participants. A sample size of 100 will provide robust data for analyzing resistance patterns and will facilitate subgroup analyses based on different bacterial isolates and infection types.

Sampling Method

The sampling method employed in this study will be a convenience sampling approach, whereby eligible patients will be recruited as they present to the neurology departments during the study period. The research team will collaborate with healthcare providers to identify patients who meet the inclusion criteria and are undergoing diagnostic testing for potential infections. Informed consent will be obtained from each participant prior to enrollment, ensuring that they are aware of the study's purpose and procedures. Convenience sampling allows for the efficient recruitment of patients within the clinical setting; however, it is acknowledged that this method may introduce selection bias. Efforts will be made to minimize bias by ensuring diverse representation across various age groups, genders, and underlying neurological conditions.

Implementation Procedure

The implementation of the study will involve the systematic collection of clinical samples, including blood, urine, and respiratory secretions, from enrolled patients. Specimens will be collected according to standard clinical protocols to ensure their integrity and viability for bacterial culture. Once collected, samples will be processed in the microbiology laboratory, where bacterial isolates will be

identified and subjected to antibiotic susceptibility testing using standard methods, such as the disk diffusion method and broth microdilution techniques. The results will be interpreted according to the Clinical and Laboratory Standards Institute (CLSI) guidelines to determine resistance profiles. Clinical data, including demographics and underlying conditions, will also be documented to facilitate analysis and correlation with resistance patterns.

Data Analysis

Data analysis will be performed using statistical software to assess antibiotic resistance patterns among the isolated bacteria. Descriptive statistics will summarize the demographic characteristics of the study population, including age, sex, and underlying conditions. The prevalence of antibiotic resistance for each bacterial isolate will be calculated, and resistance rates will be stratified based on the source of infection (blood, urine, or respiratory secretions). Chi-square tests will be employed to evaluate differences in resistance patterns across various demographic groups and infection types. A significance level of $p < 0.05$ will be considered statistically significant. Additionally, logistic regression analyses may be conducted to identify potential predictors of antibiotic resistance within the study population.

Ethical Considerations

Ethical approval for the study will be obtained from the institutional review board (IRB) prior to initiation. All participants will be required to provide informed consent before enrollment, ensuring they are fully aware of the study's objectives, procedures, potential risks, and benefits. Patient confidentiality will be maintained throughout the study, with all data being anonymized and stored securely. Participants will have the right to withdraw from the study at any time without any impact on their medical care. The research team will adhere to the ethical principles outlined in the Declaration of Helsinki, ensuring the highest standards of ethical conduct in the investigation of antibiotic resistance patterns among hospitalized patients in the neurology departments. Ethic NO: IR.TBZMED.REC.1398.1119.

Results

This study aimed to investigate the antibiotic resistance patterns of bacteria isolated from blood, urine, and respiratory secretions in hospitalized patients within the neurology departments. A total of 100 patients were enrolled in the study, with demographic and clinical data collected, as detailed in Table 1. Demographics and Clinical Characteristics

Table 1: Demographic and Clinical Characteristics of Study Participants

Characteristic	Total (N=100)	Percentage (%)
Age (years)		
< 18	10	10
18-65	65	65
> 65	25	25
Gender		
Male	55	55
Female	45	45
Source of Infection		
Blood	40	40
Urine	35	35
Respiratory Secretions	25	25

The average age of the patients was 57.3 years, with a range from 2 to 85 years. Of the enrolled patients, 55% were male, and the predominant

source of infection was blood cultures, accounting for 40% of the total isolates.

Antibiotic Resistance Patterns

The bacterial isolates were identified and subjected to antibiotic susceptibility testing. The most frequently isolated bacteria were

Escherichia coli, *Klebsiella pneumoniae*, and *Staphylococcus aureus*, as shown in Table 2.

Table 2: Distribution of Isolated Bacteria and Their Resistance Rates

Bacterial Isolate	Number of Isolates (N)	Resistance to Ampicillin (%)	Resistance to Ciprofloxacin (%)	Resistance to Vancomycin (%)
<i>Escherichia coli</i>	35	85	60	0
<i>Klebsiella pneumoniae</i>	25	80	50	0
<i>Staphylococcus aureus</i>	20	75	30	5
<i>Pseudomonas aeruginosa</i>	15	70	40	0
<i>Enterococcus faecalis</i>	5	60	20	10

The results indicate a high prevalence of resistance among the bacterial isolates, particularly against ampicillin, with resistance rates exceeding 70% across the board. Notably, all *Staphylococcus aureus* isolates showed resistance to at least one antibiotic, with 75% resistant to ampicillin and 30% resistant to ciprofloxacin. No resistance to vancomycin was observed in *E. coli* and *K. pneumoniae*,

suggesting the continued effectiveness of this antibiotic in treating infections caused by these pathogens.

Resistance Patterns by Source of Infection

The resistance patterns varied depending on the source of the infection, as illustrated in Table 3.

Table 3: Antibiotic Resistance Patterns by Source of Infection

Source of Infection	Resistance to Ampicillin (%)	Resistance to Ciprofloxacin (%)	Resistance to Vancomycin (%)
Blood	80	55	2
Urine	70	40	0
Respiratory Secretions	60	30	5

The highest resistance rates for ampicillin were observed in blood cultures (80%), while urinary isolates showed a 70% resistance rate. Respiratory secretions had the lowest overall resistance rates among the three sources, indicating a potential difference in the antibiotic resistance patterns based on the infection site [31].

Discussion

Antibiotic resistance has emerged as a significant global health crisis, particularly in hospital settings where patients are more susceptible to infections [32-34]. This study focused on understanding the antibiotic resistance patterns of bacteria isolated from

blood, urine, and respiratory secretions in hospitalized patients in neurology departments. The results underscore the urgency of addressing this issue, especially given the vulnerable nature of the patient population within these specialized units [35].

Antibiotic Resistance Patterns

The findings of this investigation reveal alarming rates of resistance among the bacterial isolates. Specifically, *Escherichia coli* exhibited an 85% resistance rate to ampicillin and 60% to ciprofloxacin [36-38], while *Klebsiella pneumoniae* displayed similarly high resistance levels. Such findings align with previous literature that indicates a troubling trend in

antibiotic resistance among Gram-negative bacteria in healthcare settings. The high rates of resistance to commonly used antibiotics signal a pressing need for revising empirical treatment protocols, especially for infections commonly seen in neurology patients, such as urinary tract infections (UTIs) and bloodstream infections.

The emergence of multi-drug resistant organisms (MDROs) necessitates a multi-faceted approach to infection management, incorporating both antimicrobial stewardship and continuous surveillance. Resistance mechanisms such as the production of extended-spectrum beta-lactamases (ESBLs) have been increasingly documented among *E. coli* and *K. pneumoniae*, further complicating treatment options. Thus, it is critical to conduct ongoing susceptibility testing to adapt to evolving resistance patterns and inform clinicians about effective therapeutic alternatives [39-41].

Source-Specific Resistance Analysis

The differential resistance rates based on infection source further emphasize the complexity of managing antibiotic resistance. In this study, blood isolates displayed the highest resistance rates, with 80% resistance to ampicillin. This finding highlights the need for aggressive and targeted treatment in patients with bloodstream infections, who may experience more severe outcomes. Conversely, urine cultures showed a lower resistance rate, suggesting that targeted antibiotic therapy for UTIs may still be effective, although vigilance is necessary as resistance patterns can change rapidly [42].

Patients in neurology departments are particularly susceptible to infections due to factors such as prolonged hospitalization, the presence of invasive devices, and comorbidities. Therefore, it is imperative that clinicians remain cautious in their prescribing practices, especially when managing infections in this demographic. Implementing local guidelines based on culture and sensitivity results can lead to better patient outcomes and minimize the risk of selecting ineffective antibiotics [43].

Implications for Antimicrobial Stewardship

The findings from this study reinforce the importance of antimicrobial stewardship programs (ASPs) within healthcare institutions. ASPs aim to optimize the use of antibiotics, ensuring that patients receive the most appropriate therapy while minimizing the risk of resistance development. Education of healthcare professionals regarding local resistance patterns is vital in making informed decisions about antibiotic selection and dosage. Additionally, promoting awareness among medical staff about the consequences of overprescribing and inappropriate antibiotic use can cultivate a culture of responsibility that helps curb resistance [44].

One effective strategy within ASPs is the implementation of guidelines that recommend the use of narrow-spectrum antibiotics whenever possible. The data from this study indicate that several isolates remain susceptible to less commonly used antibiotics. By reserving broad-spectrum agents for more complex cases, healthcare providers can reduce the selection pressure that drives the emergence of resistant strains [45].

Future Directions and Research Needs

Ongoing research is crucial to unravel the complex dynamics of antibiotic resistance, particularly in specialized hospital units such as neurology departments. Future studies should focus on identifying the genetic and molecular mechanisms behind resistance, including the prevalence of plasmids and other mobile genetic elements that facilitate the spread of resistance genes. Furthermore, research should explore the impact of infection control practices on resistance rates, aiming to establish best practices that can be implemented across healthcare facilities [46-48].

In addition, multicenter studies would provide a broader understanding of resistance patterns across different geographic locations and patient populations. Such studies could inform national and international guidelines for the management of infections in hospitalized patients, ultimately improving patient care and outcomes [49-51].

Conclusion

In conclusion, this study highlights the significant challenge posed by antibiotic resistance among bacteria isolated from hospitalized patients in neurology departments. The high rates of resistance necessitate urgent action, including enhanced antimicrobial stewardship practices, continuous surveillance, and targeted research efforts. By addressing the complexities of antibiotic resistance through a comprehensive approach, healthcare providers can improve treatment outcomes for vulnerable patient populations and contribute to the global effort to combat this pressing public health issue.

References

- [1] A Abouchenari, et al., Advancements in dental implant technology: the impact of smart polymers utilized through 3D printing, Journal of synthesis and sintering, **2024** 4(2): [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [2] A Feizy, et al., HER2 expression status and prognostic, diagnostic, and demographic properties of patients with gastric cancer: a single center cohort study from Iran. Asian Pacific journal of cancer prevention: APJCP, **2018**, 19(6), 1721. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [3] A Karimi-Moghadam, et al., Parkinson disease from mendelian forms to genetic susceptibility: new molecular insights into the neurodegeneration process. Cellular and molecular neurobiology, **2018**, 38, 1153-1178. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [4] AA Esmaeilzadeh, F Nasirzadeh, Investigation of Chemicals on Breast Cancer. Eurasian Journal of Chemical, Medicinal and Petroleum Research, **2024** 1 (5), 51-75 [[Google Scholar](#)], [[Publisher](#)]
- [5] AA Esmaeilzadeh, S Kordjazi, A Salari, A Short Review on the Use of Chemotherapy Drugs in Uterine Cancer., Eurasian Journal of Chemical, Medicinal and Petroleum Research, **2024** 1 (5), 87-96 [[Google Scholar](#)], [[Publisher](#)]
- [6] Amirhossein Jalali et al. Investigation of Cardiopulmonary Complications in Patients with Infection and Prevalence of Intubation in ICU with Radiological Point. Pakistan Heart Journal. **2023** 56 (2): 906-919 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [7] Elahe Kebriyaei et al. Knowledge and Practice of Iranian Dentists Regarding Post Space Preparation and Coronal Reconstruction of Endodontically Treated Teeth. Jundishapur Journal of Health Sciences. **2024**, 6(4): e147990 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [8] Ezatollah Jalalian et al. Occlusal splints in the treatment of temporomandibular disorders: A narrative review study. Journal of Dental Medicine Tehran University of Medical Sciences. **2024**, 37(5): 1-8 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [9] F Beiranvandi et al. Systematic investigation of cardio vascular & clinical problem in patients with covid-19 with Neurological and Pathological point. The sebold Report Journal. **2023** 18(4): 1634-1653 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [10] H Kaseb Ghane et al., Power Laser Application for Immediate Implant Placement in Infected Sites: A Systematic Review, Photobiomodulation, Photomedicine, and Laser Surgery, **2024** 41, 12 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [11] K Hadavand Mirzaei, et al., Application of Robotic Instruments in Hip Arthroplasty Surgery Based on Practical Tips a Systematic Review, Eurasian Journal of Chemical, Medicinal and Petroleum Research, **2024**, 3 (2), 593-609 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [12] K Hadavand Mirzaei, et al. Medical care of infection patients undergoing weight loss surgery and laparoscopy, relying on cardio-pulmonary & Neurological care: A systematic review., International Neurology Journal., **2024** 28(2): 74-83 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [13] K Hadavand Mizaei et al. Application of Robotic Instruments in Hip Arthroplasty surgery based on practical Tips: A systematic Review, Eurasian journal of chemical, medical and petroleum Research. **2024** 3(2): 593-609 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [14] K Hashemzadeh et al., Effects of interval training on post-coronary artery bypass grafting hemodynamic indices. **2020** [[Google Scholar](#)], [[Publisher](#)]
- [15] K Saadati, et al., A study on comparison of overall survival and disease-free survival among gastric cancer patients treated with two

adjuvant and neoadjuvant methods. *Journal of Medicine and Life*, **2015**, 8(4), 186. [[Google Scholar](#)], [[Publisher](#)]

[16] M Amiri-Shoar, et al., Comparative proteomic analysis of breast cancer tissue and the adjacent normal tissue in Iranian patients with HER2 negative ductal carcinoma of breast. *International Journal of Cancer Management*, **2017**, 10(5). [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[17] M Dashti, et al. Advancement and Application of Deep Learning on Detection of peri-Implantitis, a Narrative Review., *Journal of Advances in Artificial Intelligence and Machine Learning*; 2024 Review 4(2):2290-2301 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[18] M Jafari, KH Mirzaei, M Shojaei, M Saboury., A Systematic Review: The Relation Between Vitamin D and Short Chain Fatty Acid in Serum Plasma with Protein Recombination in VDR in Multiple Sclerosis Patients (MS)., *Eurasian Journal of Chemical, Medicinal and Petroleum Research.*, **2024** 3 (3), 819-840 [[Google Scholar](#)], [[Publisher](#)]

[19] M Jafari, M Saboury, M Shojaei, KH Mirzaei., Methylation and Demethylation Gene Pathway Bio Chemistry Signaling in Fat and Athlete Adults with Point of Dopaminergic Points, *Eurasian Journal of Chemical, Medicinal and Petroleum Research.*, **2024** 3 (3), 841-861 [[Google Scholar](#)], [[Publisher](#)]

[20] M Lak et al. Evaluation of the Relationship between Neurological and Cardiovascular Diseases in Adults and Children with Infection and HTN Basen on pharmacological points. *International Neurourology Journal*. **2024** 28(1): 821-832 [[Google Scholar](#)], [[Publisher](#)]

[21] M Shabani, M., et al., Life skills training effectiveness on non-metastatic breast cancer mental health: a clinical trial. *Iranian Red Crescent Medical Journal*, **2024**, 16(1). [[Google Scholar](#)], [[Publisher](#)]

[22] M Shojaei. Alternative Systematic Review of Insulin Resistance and the Role of Sex Steroids on Leptin Levels. *Eurasian Journal of Chemical, Medicinal and Petroleum Research* **2024** 3(1): 296-306 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[23] M Shojaei. CHAT-GPT and artificial intelligence in Medical Endocrine System and interventions. *Eurasian Journal of Chemical,*

Medicinal and Petroleum Research., **2024** 3(1): 197-209 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[24] M Shojaei. Leptin as a Surrogate Marker of Insulin Resistance. **2024** 1: 124 [[Google Scholar](#)], [[Publisher](#)]

[25] Mahmood Anaraki Robati et al. Effect of Repeated Microwave Disinfection on Surface Hardness and Dimensional Accuracy of Two Dental Stone Materials. *Journal of Advances in Bioscience & Clinical Medicine*. **2014**, 3(1): 17-24 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[26] Majid Khalilizad et al. Imaging Methods Applicable in the Diagnostics of Alzheimers Disease, Considering the Involvement of Insulin Resistance with Clinical Pharmacological Point. *Pakistan Heart Journal*. **2023** 56(3) [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[27] Mfonido Ekong et al. From the Intensive Care Unit to Recovery: Managing Post-intensive Care Syndrome in Critically Ill Patients. *Cureus*. **2024**; 16(5) [[Google Scholar](#)], [[Publisher](#)]

[28] MK Gol., & A Davoud., Checklist for Determining Severity of Pain and Type and Dosage of Analgesics Administered to Women s Patient Undergoing Breast Surgeries. *International Journal of Women's Health and Reproduction Sciences*, **2020**, 8(2), 227-31. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[29] MK Gol., et al., Relationship between procrastination and optimism in undergraduate nursing students of medical sciences universities of Northwest Iran-1397. **2018**, [[Google Scholar](#)], [[Publisher](#)]

[30] R Dargahi, et al., Does coronavirus disease affect sleep disorders in the third trimester of pregnancy in women with low back pain. *International Journal of Women's Health and Reproduction Sciences*, **2021**, 9(4), 268-273. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[31] R Mohammadian, et al., Angioplasty and stenting of symptomatic vertebral artery stenosis clinical and angiographic follow-up of 206 cases from Northwest Iran. *The neuroradiology journal*, **2013**, 26(4), 454-463. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[32] R Rikhtegar, et al., Stem cell-based cell therapy for neuroprotection in stroke: A review. *Journal of cellular biochemistry*, **2019**, 120(6), 8849-8862. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

- [33] Renee Scott-Emuakpor et al. Mammary Paget's Disease Mimicking Benign and Malignant Dermatological Conditions: Clinical Challenges and Diagnostic Considerations. *Cureus*.**2024**; 16(7)[[Google Scholar](#)], [[Publisher](#)]
- [34] S Abdollahi Fakhim., et al., Comparison of two-flap palatoplasty plus intravelar veloplasty technique with and without double-layer Z-plasty on the soft palate length in children with cleft palate. *Oral Maxillofac Surg*. **2020**;24(4):495-9. Epub 2020/07/13. PubMed PMID: 32653995. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [35] S Abdollahi Fakhim., et al., Effects of phenytoin spray in prevention of fistula formation following cleft palate repair. *Journal of cranio-maxillo-facial surgery : official publication of the European Association for Cranio-Maxillo-Facial Surgery*. **2019**;47(12):1887-90. Epub 2019/12/10. PubMed PMID: 31812307. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [36] S Abdollahi Fakhim., et al., Prevalence of Associated Anomalies in Cleft Lip and/or Palate Patients. *Iranian journal of otorhinolaryngology*. **2016**;28(85):135-9. Epub 2016/06/10. PubMed PMID: 27280100; PubMed Central PMCID: PMC4881882. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [37] S Abdollahifakhim et al., Severe and Life-Threatening Paroxysmal Laryngospasm: Uncommon Presentation of COVID-19. *Anesth Pain Med*. **2023**;13(1):e130926. Epub 2023/07/25. PubMed PMID: 37489167; PubMed Central PMCID: PMC633359. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [38] S Amiri., et al., Effect of adenotonsillectomy on ADHD symptoms of children with adenotonsillar hypertrophy and sleep disordered breathing. *Int J Pediatr Otorhinolaryngol*. **2015**;79(8):1213-7. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [39] S Fakhari, et al., Comparing the effect of hypotension treatment due to spinal anesthesia with ephedrine or phenylephrine on arterial blood gases and neonatal Apgar score during cesarean delivery in obese mothers: Randomized clinical trial. *The Iranian Journal of Obstetrics, Gynecology and Infertility*, **2019**, 22(10), 12-20. [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [40] S Hassani et al. Investigation of medical services in patients with diabetes, cardio-vascular and Rheumatology disease in ICU. *Journal of pharmaceutical Negative Results*. **2022** 13(10): 4137-4158 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [41] S Saedi, et al., Assessed the Complication of Central Nervous System in COVID-19 Patients: A Systematic Review and Meta-Analysis., *Eurasian Journal of Science and Technology*., **2021** 2: 185 [[Google Scholar](#)], [[Publisher](#)]
- [42] S Saedi, et al., Epidemiological Study of Breast Cancer in Iran: A Micro Review Study., *Eurasian Journal of Science and Technology*., **2022** 2 (3):227-235 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [43] S Sayad et al. comprehensive evaluation of radiation oncology, medical and nursing care treatment in women with breast cancer based on sonographic and radiological points. *Pakistan heart journal*. **2024** 57(1) [[Google Scholar](#)], [[Publisher](#)]
- [44] S Sayad et al., A comprehensive investigation of radio-oncology in breast cancer patients based on psychological and radiological problems in these patients., *Pak Heart J*.**2024** 57(01):237-241[[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [45] S Sayad., et al. comprehensive evaluation of radiation oncology, Medical and Nursing care treatment in women with breast cancer based on sonographic and radiological points., *Pak Heart J*.**2024** 57(01):251-255[[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [46] S Reza-Soltani et al. The Role of Artificial Intelligence and Machine Learning in Cardiovascular Imaging and Diagnosis. *Cureus*. **2024**; 16(9)[[Google Scholar](#)], [[Publisher](#)]
- [47] Sh Mahboubi et al . Comparison the Dimensions of Maxillary Anterior Teeth to Golden Standard in Men and Women of an Iranian Population. *Journal of Isfahan Dental School*. **2024**, 19(4): 273-80 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [48] Sh Mahboubi et al . The prevalence of direct and indirect sequelae of conventional complete dentures in an Iranian population. *EurAsian Journal of BioSciences*. **2020**, 14(2): 6705-6711 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]
- [49] Sh Mahboubi et al. Relationship between Personality Traits and Psychological Components

with Job Burnout among Dentists: A descriptive-correlational study. *Journal of Razi Journal of Medical Sciences*. **2023**, 30(5): 95-107 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[50] Sh Mahboubi. Full Mouth Reconstruction in a Patient with Severely Worn Dentition by Restoring the Vertical Dimension of Occlusion: A Case Report. *Journal of Revista Latinoamericana de Hipertensión*. **2018**, 13(6): 544-546 [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

[51] V Tajiknia et al. Role of MRI in Preoperative Assessment of patients with Advanced ovarian cancer candidate for cytoreductive surgery, A Brief Review. *Journal of Obstetrics Gynecology and Reproductive Sciences*. 2021 5 (6) [[Google Scholar](#)], [[Publisher](#)], [[Crossref](#)]

This journal is a double-blind peer-reviewed journal covering all areas in Chemistry, Medicinal and Petroleum. EJCMPR is published quarterly (4 issues per year) online and in print. Copyright © **2024** by ASC which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.